



# Effectiveness of intravenous immunoglobulin for children with severe COVID-19: a rapid review

Jingyi Zhang<sup>1,2#</sup>, Yinmei Yang<sup>3,4,5#</sup>, Nan Yang<sup>2#</sup>, Yanfang Ma<sup>2</sup>, Qi Zhou<sup>6</sup>, Weiguo Li<sup>3,4,5</sup>, Xia Wang<sup>3,4,5</sup>, Liping Huang<sup>3,4,5</sup>, Xufei Luo<sup>1,2</sup>, Toshio Fukuoka<sup>7,8</sup>, Hyeong Sik Ahn<sup>9,10</sup>, Myeong Soo Lee<sup>11,12</sup>, Zhengxiu Luo<sup>3,4,5</sup>, Yaolong Chen<sup>1,2,13,14,15</sup>, Enmei Liu<sup>3,4,5</sup>, Kehu Yang<sup>1,2,15</sup>, Zhou Fu<sup>3,4,5</sup>, on behalf of COVID-19 Evidence and Recommendations Working Group

<sup>1</sup>School of Public Health, Lanzhou University, Lanzhou 730000, China; <sup>2</sup>Evidence-based Medicine Center, School of Basic Medical Sciences, Lanzhou University, Lanzhou 730000, China; <sup>3</sup>Department of Respiratory Medicine, Children's Hospital of Chongqing Medical University, Chongqing 400014, China; <sup>4</sup>National Clinical Research Center for Child Health and Disorders, Ministry of Education Key Laboratory of Child Development and Disorders, China International Science and Technology Cooperation Base of Child Development and Critical Disorders, Children's Hospital of Chongqing Medical University, Chongqing 400014, China; <sup>5</sup>Chongqing Key Laboratory of Pediatrics, Chongqing 400014, China; <sup>6</sup>The First School of Clinical Medicine, Lanzhou University, Lanzhou 730000, China; <sup>7</sup>Emergency and Critical Care Center, the Department of General Medicine, Department of Research and Medical Education at Kurashiki Central Hospital, Okayama, Japan; <sup>8</sup>Advisory Committee in Cochrane Japan, Tokyo, Japan; <sup>9</sup>Department of Preventive Medicine, Korea University College of Medicine, Seoul, Korea; <sup>10</sup>Korea Cochrane Centre, Seoul, Korea; <sup>11</sup>Korea Institute of Oriental Medicine, Daejeon, Korea; <sup>12</sup>University of Science and Technology, Daejeon, Korea; <sup>13</sup>Lanzhou University, an Affiliate of the Cochrane China Network, Lanzhou 730000, China; <sup>14</sup>Chinese GRADE Center, Lanzhou 730000, China; <sup>15</sup>Key Laboratory of Evidence Based Medicine and Knowledge Translation of Gansu Province, Lanzhou University, Lanzhou 730000, China

**Contributions:** (I) Conception and design: Y Chen, E Liu; (II) Administrative support: K Yang, Z Fu; (III) Provision of study materials or patients: J Zhang, Y Yang; (IV) Collection and assembly of data: Y Yang, N Yang; (V) Data analysis and interpretation: Y Ma, J Zhang; (VI) Manuscript writing: All authors; (VII) Final approval of manuscript: All authors.

#These authors contributed equally to this work.

**Correspondence to:** Zhou Fu. Department of Respiratory Medicine, Children's Hospital of Chongqing Medical University, Chongqing 400014, China. Email: fu\_zhou79@aliyun.com; Kehu Yang. School of Public Health, Lanzhou University, Lanzhou 730000, China. Email: kehuyangebm2006@126.com.

**Background:** Intravenous immunoglobulin (IVIG) is usually used as supportive therapy, but the treatment of COVID-19 by IVIG is controversial. This rapid review aims to explore the clinical effectiveness and safety of IVIG in the treatment of children with severe COVID-19.

**Methods:** We systematically searched the literature on the use of IVIG in patients with COVID-19, severe acute respiratory syndrome (SARS) or Middle East respiratory syndrome (MERS), including both adults and children. We assessed the risk of bias and quality of evidence and reported the main findings descriptively.

**Results:** A total of 1,519 articles were identified by initial literature search, and finally six studies met our inclusion criteria, included one randomized controlled trial (RCT), four case series and one case report involving 198 patients. One case series showed the survival of COVID-19 patients with acute respiratory distress syndrome (ARDS) was not improved by IVIG. One case report showed high-dose IVIG could improve the outcome of COVID-19 adults. Three observational studies showed inconsistent results of the effect of IVIG on SARS patients. One RCT showed that IVIG did not reduce mortality or the incidence of nosocomial infection in adults with severe SARS. The quality of evidence was between low and very low.

**Conclusions:** The existing evidence is insufficient to support the efficacy or safety of IVIG in the treatment of COVID-19.

**Keywords:** COVID-19; children; intravenous immunoglobulin (IVIG); rapid review

Submitted Apr 14, 2020. Accepted for publication Apr 26, 2020.

doi: 10.21037/atm-20-3305

View this article at: <http://dx.doi.org/10.21037/atm-20-3305>

## Introduction

COVID-19 is an infectious disease caused by severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2) (1). SARS-CoV-2 belongs to the family of coronaviruses, which are enveloped viruses that can cause illnesses ranging from common cold to severe diseases such as SARS and MERS (2,3). The COVID-19 epidemic massively influences the public health and people's daily lives, and the disease was declared a pandemic on 11 March 2020 (4). All populations are susceptible to infection and there is a research shows children are as likely to be infected as adults (5). On February 11, 2020, there were 44,672 confirmed cases in mainland China, of whom 416 were under the age of 10 years and 549 between the ages of 10 to 19 years (6). The main symptoms in children are fever and cough, and the disease is on average less severe in children than adults (7). However, severe cases have been reported also in children (8). So far, there has been no specific treatment for COVID-19, antiviral therapy and vaccination are currently under development (9,10).

IVIG is prepared from the plasma of healthy humans and usually used as supportive therapy. Its main component is immunoglobulin (Ig) G, which has dual therapeutic effects of immune-modulation effects and immune substitution (11). IVIG is one of the alternative treatments for children with agammaglobulinemia, and an effective treatment of Kawasaki disease (12,13). IVIG was used to treat SARS patients during the SARS outbreak in 2003 (14,15), but there is no convincing evidence of its effectiveness. According to recent reports, about 33% of patients with severe COVID-19 received IVIG in China (16). Some published guidelines of COVID-19 have indicated that IVIG could be used to treat children with severe or critical disease (17).

The purpose of this study is to perform a comprehensive rapid review to explore whether it is beneficial to treat children with severe COVID-19 with IVIG and provide supporting evidence support for COVID-19 guidelines. Because of the urgent situation, the review was not registered (18). We present the following article in accordance with the PRISMA reporting checklist (available at <http://dx.doi.org/10.21037/atm-20-3305>).

## Methods

### Search strategy

We carried out a comprehensive search in the following electronic databases: the Cochrane library, MEDLINE

(via PubMed), EMBASE, Web of Science, China Biology Medicine disc (CBM), China National Knowledge Infrastructure (CNKI), and Wanfang Data, by using the terms “COVID-19”, “SARS-CoV-2”, “Novel coronavirus”, “2019-novel coronavirus”, “2019-nCoV”, “SARS”, “MERS”, “IVIG”, “intravenous immunoglobulin” and their derivatives. The search covered the time from each database's inception to March 31, 2020. The search strategies were determined by multiple pre-searches and were discussed with the clinicians about the appellation of disease and IVIG. We also searched the World Health Organization Clinical Trials Registry Platform, ISRCTN Registry, ClinicalTrials, Google Scholar, three preprint services, including medRxiv (<https://www.medrxiv.org/>), bioRxiv (<https://www.biorxiv.org/>) and SSRN (<https://www.ssrn.com/index.cfm/en/>) and references of included studies. The details of the search strategy can be found in the *supplementary material*.

### Inclusion and exclusion criteria

We included RCTs that compared IVIG treatment (standard intravenous immunoglobulin (IVIG) preparations, excluding IgM-enriched Ig, hyperimmune Ig and specific Ig from convalescent plasma) with a control group (placebo or no treatment with IVIG), and cohort studies, cross-sectional studies, case-control studies, case series and cases report that can distinguish the corresponding outcomes caused by IVIG. The inclusion of COVID-19 adult patients and patients with SARS or MERS helps to provide indirect evidence, if studies on COVID-19 in children are scarce. Studies with all patients diagnosed with COVID-19, SARS or MERS were included, without restrictions on age, race, gender, or geographical location or setting. The primary outcomes were the risk of death and survival probability. Secondary outcomes included the incidence of nosocomial infection, the duration of fever, time of the lung lesions subsided obviously, the progression of disease cascade, improvement in abnormal laboratory inspection indicators (total peripheral blood WBC, platelet counts, serum globulin, WBC counts), and adverse effects. We excluded duplicates, conference abstracts, comments and letters, studies published in languages other than English or Chinese, and studies where we could not access the full text.

### Study selection

After eliminating duplicates by EndNote software and

manual check, two reviewers (J Zhang and Y Yang) independently reviewed the titles and abstracts of records retrieved from the search and selected all potentially relevant studies according to the pre-defined inclusion and exclusion criteria. After this, the same reviewers screened the full texts and made the final selection. A pilot search was conducted before the full screening of the literature to ensure that each researcher understood the screening criteria and process. Disagreements about selection of studies were resolved by consulting a third reviewer (N Yang). The process of study selection was documented using a PRISMA flow diagram (19).

### **Data extraction**

Two reviewers (J Zhang and Y Yang) independently extracted the following data from included trials using a standardized extraction sheet: (I) basic information (year of publication, first author and affiliation, journal, funding, conflict of interest); (II) study details (type of study, sample size, research purpose, research population characteristics, interventions; and (III) outcome data. A pre-test was conducted before formal extraction to ensure that each researcher agreed with the extraction criteria and process. Disagreements were solved through discussion with a third reviewer (N Yang).

### **Risk of bias assessment**

Two reviewers (J Zhang and Y Ma) assessed the quality of the included studies independently. We used the Cochrane bias risk assessment tool (Risk of bias) to assess RCTs (20), the criteria recommended by the National Institute of Health and Clinical Optimization (NICE) for case series to assess the risk of bias (21), the Joanna Briggs Institute's (JBI) case report quality appraisal tool for case reports (22) Newcastle-Ottawa Quality Assessment Scale (NOS) for the quality of cohort studies and case-control studies (23), and Agency for Healthcare Research and Quality (AHRQ) tool for cross-sectional studies (24).

### **Data synthesis**

If the data were enough to be summarized, we would conduct a meta-analysis using Review Manager 5.3. For dichotomous outcomes we calculated the risk ratio (RR) and the corresponding 95% confidence interval (CI) and

P value. For continuous outcomes, we calculated the mean difference (MD) and its corresponding 95% CI when means and standard deviations (SD) were reported. If sufficient data were available, we considered examining the robustness of meta-analyses in a sensitivity analysis. When effect sizes could not be pooled, we reported the study effectiveness narratively.

### **Quality of the evidence assessment**

The quality of the body of evidence was graded using the GRADE method (25,26). Evidence from randomized trials could be downgraded by the following five factors: risk of bias, inconsistency of results, indirectness of evidence, imprecision of results, and publication bias (*Table S1*). The quality of evidence for each outcome was graded as high, medium, low, or very low. The results of the grading were presented in "GRADE evidence profile" (27-30).


## **Results**

We identified 1,519 articles in the initial literature search (*Figure 1*). After removing duplicates, we screened the titles and abstracts of 1,405 records. Thirty-one articles were retrieved for full-text reviewing. Finally, one RCT, four case series and one case report involving a total of 198 patients were included for rapid review (31-36).

The studies were published between 2003 and 2020, and all studies were from China (*Table 1*). We found one case series on IVIG in COVID-19 adults with acute respiratory distress syndrome (ARDS), one case report in COVID-19 adults, one RCT of 44 adults with severe SARS, which included 25 adults with acute lung injury (ALI) and 19 adults with ARDS, one case series involving children with SARS, and two case series also involving adults with SARS. In all studies IVIG was used before or in combination with other drugs and treatment (such as antibiotics, glucocorticoids, antivirals, oxygen therapy). The IVIG dose and duration of use differed across studies.

### **Risk of bias for included studies**

We found a high risk of bias in random sequencing, allocation concealment and blinding in the only included RCT. All case series had a moderate risk (score 4 to 5 out of 8) (*Table 1*), one case report meeting 8 of the 8 items of the JBI quality appraisal tool.



**Figure 1** The flow chart of the literature search.

**Table 1** Basic characteristics of the included studies

Author	Study location	Study design	Number (M/F)	Disease	Age (range or mean $\pm$ SD)	Intervention		Outcome	Risk of bias <sup>#</sup>
						Other treatment*	IVIG		
Zeng 2003 (33)	Guangzhou	Case series	5/5	SARS	7.3 $\pm$ 5.1	Antibiotics, oxygen inhalation, symptomatic, comprehensive treatment, etc.	200–400 mg/kg/d for 3 days	I, II, III	4/8
Wu 2003 (35)	Guangzhou	Case series	66	SARS	16–62	Antibiotics, glucocorticoids, interferons, antivirals and oxygen therapy	5–10 g/d for 3 to 6 days	IV	4/8
Wang 2004 (34)	Taiwan	Case series	22	SARS	24–87	Methylprednisolone	1 g/kg/d for 2 days	V, VI	5/8
Wu 2005* (36)	Guangzhou	RCT	15/29	SARS	Mean: 42/43	Antibiotics, glucocorticoids	2.5 mg/d for 2 days–10 mg/d for 13 days	VII, VIII, IX	NA
Liu 2020 (31)	Wuhan	Case series	29/24	COVID-19	Mean: 55	Antibiotics, glucocorticoids, interferons, antivirals and oxygen therapy	NR	X	5/8
Cao 2020 (32)	Wuhan	Case report	3	COVID-19	34–56	NR	25 g/d for 5 days	XI	NA

Outcome: I: the duration of fever; II: total peripheral blood WBC ( $10^9/L$ ); III: time of the lung lesions subsided obviously; IV: adverse effects; V: WBC counts ( $10^9/L$ ); VI: platelet counts ( $10^9/L$ ); VII: serum globulin (g/L); VIII: the incidence of nosocomial infection; IX: the risk of death; X: survival probability; XI: the progression of disease cascade. \*, prior to IVIG treatment, the patient received other treatment; #, risk of bias in case series. NR, not report; NA, not applicable.

### Quality of the evidence

The quality of evidence for all outcomes assessed in the only included RCT was graded low (*Table S2*), primarily due to serious risk of bias and imprecision. As we included four case series, we judged that reporting a 'GRADE evidence profile' would not be meaningful. Overall, the quality of evidence was very low for most outcomes and cannot thus provide a reliable indication of any likely effect across outcomes.

### COVID-19

A case series of 109 adults with COVID-19 reported that most patients used antibiotics and antiviral treatment, and over half of the patients were given glucocorticoid therapy and IVIG. The survival probability of patients with ARDS could not be improved by antiviral, glucocorticoid, or Ig treatment. The risk of death was not associated with the use of IVIG in the patients with ARDS (31). A case report of three adults with COVID-19 reported that a high dose IVIG (25 g/d for 5 days) administered at the appropriate point could successfully block the progression of disease cascade (result of the clinical symptoms, laboratory inspection indicators and chest CT scan), and finally improve the outcome of COVID 19 (32).

### SARS

#### The incidence of nosocomial infection and the risk of death

The RCT of 44 adults with severe SARS found no significant difference in the risk of death (18.1% *vs.* 23.8%) or the incidence of nosocomial infection (65.2% *vs.* 52.4%) between adults treated either with IVIG or with conventional treatment. And there was no significant difference in the incidence of nosocomial infection between ALI (50.0% *vs.* 38.5%) and ARDS (81.8% *vs.* 75.0%) patients (36).

#### Laboratory inspection

One case series reported the patients with SARS who did not receive steroids for severe hemocytopenia had increased WBC counts and platelet counts after undergoing IVIG (34). Another case series reported the children with persistent fever who were given IVIG had significantly improved total peripheral blood WBC after undergoing

IVIG (33). The included RCT showed the serum globulin increased slightly in the IVIG group, but decreased in the conventional treatment group and the difference was not significant (36).

#### The duration of fever

One case series included ten children with persistent fever who were given IVIG. The body temperature ranged between 38.4 and 40 °C at baseline and the duration of fever was 1 to 4 days after IVIG (33).

#### Imaging testing

One case series reported that chest radiographs in children who were given IVIG showed more patchy focal asymmetric infiltrative shadows, more rapid time of the lung lesions subsided obviously than in a randomly selected, age- and sex-matched control group of 20 children without IVIG (33).

#### Adverse effects

One case series reported no adverse effects associated with the use of IVIG in the early stage of the disease in patients with high fever and other symptoms, or for patients with WBC count below  $3.0 \times 10^9/L$  (35).

### Discussion

We only found limited evidence about the use of IVIG to treat children or adults with severe COVID-19. Since SARS and COVID-19 belong to the same family of viruses, we used IVIG treatment of SARS as indirect evidence, even though the quality of the included studies was generally low. The results were also inconsistent, and no benefit was found in the only identified RCT.

An earlier systematic review of treatment effects with SARS concluded that although four studies suggested an improvement in the patients' condition after IVIG treatment, more controlled trials are needed to provide evidence of the potential benefits on IVIG against SARS. The results of the review are roughly in line with our findings, more high-quality evidence about the benefits and disadvantages of IVIG for COVID-19 and SARS are needed (37).

There was no apparent benefit from IVIG, despite it being used to treat other respiratory infections. A meta-analysis of seven RCTs in children aged less than three years with respiratory syncytial virus infection found no evidence of differences between children treated with IVIG or with

placebo in the risk of death (RR =0.87, 95% CI: 0.14–5.27) or serious adverse events (RR =1.08, 95% CI: 0.65–1.79), or in the duration of hospitalization (MD =-0.70, 95% CI: -1.83 to 0.42) (38). SARS belongs to the category of systemic inflammatory response syndromes (SIRS), and severe SARS often manifests as ALI, ARDS and progresses to severe sepsis (39,40). The course of COVID-19 may also be similar. A meta-analysis of nine RCTs showed that IVIG did not reduce the mortality (OR =0.95, 95% CI: 0.80–1.13), length of hospital stay (MD =-4.08, 95% CI: -6.47 to -1.69), or the risk of death or major disability before two years of age (RR =0.98, 95% CI: 0.88–1.09) in infants with suspected or confirmed infection, compared with placebo or no intervention (41).

IVIG is prepared from pools of plasma obtained from several thousand healthy blood donors. Unlike convalescent plasma from patients with COVID-19, IVIG does not contain SARS-Cov-2 neutralizing antibody (42,43). The review showed that there was no evidence that IVIG has an effect on anti-MERS-CoV, or that IVIG would cause kidney failure or thrombosis in patients with MERS (44). IVIG could increase the risk of vaccination delay. A study by National Advisory Committee on Immunization and the American Advisory Committee on Immunization showed a delay of five months of varicella vaccine in patients who received IVIG and varicella immune globulin (VZIG) (45). IVIG may increase the risk of infections transmitted by transfusion (42). Some adverse effects, such as thrombosis, aseptic meningitis, hemolysis, and renal failure, are mainly associated with the use of high-dose IVIG (46).

The condition, dose and duration of IVIG were inconsistent between studies, the efficacy, and the associated adverse effects remain unclear. The first severe case of COVID-19 in children in China took IVIG with a dose of 400 mg/kg for a duration of five days (8). The recommended dosage of IVIG for children with severe COVID-19 was also inconsistent in different guidelines, including 1.0 g/kg/d for 2 days, or 400 mg/kg/d for 5 days, 0.2 g/kg/d for 3–5 days, or 1–2 g/kg for 2–3 days (47–49).

Therefore, it is particularly important and urgent to study the benefits and disadvantages of IVIG treatment in children with COVID-19. It is promising that a trial addressing efficacy and safety of IVIG therapy in patients with severe or critical COVID-19 disease has been registered on (50). Randomized, double-blinded, large sample, multicenter clinical trials on children are urgently

needed for getting scientific evidence to support clinical decision-making.

### ***Strength and limitations***

This is the first rapid review of IVIG treatment for children with COVID-19. There are several limitations in this systematic review. First, the use of glucocorticoids or a combination of a variety of broad-spectrum antibiotics before IVIG may lead to changes in the microecology of the body, affect the immune regulation function, and thus also affect the effect of IVIG. Second, the total sample size of this study was insufficient to make strong conclusions, and the quality of the methodology was generally low which affect the certainty of the results. Finally, we may have missed some studies as we only included studies published in Chinese and English.

### **Conclusions**

There is no direct evidence for IVIG in children with COVID-19, current evidence is insufficient to assess the effectiveness and safety of IVIG for children with severe COVID-19. Therefore, we cannot suggest use of IVIG for the treatment of COVID-19 in children. More clinical studies to address this topic are needed.

### **Acknowledgments**

We thank Janne Estill, Institute of Global Health of University of Geneva for providing guidance and comments for our review. We thank all the authors for their wonderful collaboration.

**Funding:** This work was supported by grants from National Clinical Research Center for Child Health and Disorders (Children's Hospital of Chongqing Medical University, Chongqing, China) (grant number NCRCCHD-2020-EP-01); Special Fund for Key Research and Development Projects in Gansu Province in 2020; The fourth batch of "Special Project of Science and Technology for Emergency Response to COVID-19" of Chongqing Science and Technology Bureau. Special funding for prevention and control of emergency of COVID-19 from Key Laboratory of Evidence Based Medicine and Knowledge Translation of Gansu Province (grant number No. GSEBMKT-2020YJ01); The Fundamental Research Funds for the

Central Universities (lzujbky-2020-sp14).

## Footnote

*Reporting Checklist:* The authors have completed the PRISMA reporting checklist. Available at <http://dx.doi.org/10.21037/atm-20-3305>

*Conflicts of Interest:* All authors have completed the ICMJE uniform disclosure form (available at <http://dx.doi.org/10.21037/atm-20-3305>). MSL serves as the unpaid editorial board member of *Annals of Translational Medicine* from Nov 2019 to Oct 2021. The other authors have no conflicts of interest to declare.

*Ethical Statement:* The authors are accountable for all aspects of the work in ensuring that questions related to the accuracy or integrity of any part of the work are appropriately investigated and resolved.

*Open Access Statement:* This is an Open Access article distributed in accordance with the Creative Commons Attribution-NonCommercial-NoDerivs 4.0 International License (CC BY-NC-ND 4.0), which permits the non-commercial replication and distribution of the article with the strict proviso that no changes or edits are made and the original work is properly cited (including links to both the formal publication through the relevant DOI and the license). See: <https://creativecommons.org/licenses/by-nc-nd/4.0/>.

## References

- Zhu N, Zhang D, Wang W, et al. A novel coronavirus from patients with pneumonia in China, 2019. *N Engl J Med* 2020;382:727-33.
- Chan JFW, Kok KH, Zhu Z, et al. Genomic characterization of the 2019 novel human-pathogenic coronavirus isolated from a patient with atypical pneumonia after visiting Wuhan. *Emerg Microbes Infect* 2020;9:221-36.
- Lu R, Zhao X, Li J, et al. Genomic characterisation and epidemiology of 2019 novel coronavirus: implications for virus origins and receptor binding. *Lancet* 2020;395:565-74.
- The World Health Organization, Coronavirus disease (COVID-19) Pandemic[Internet]. World Health Organization; c2020 [cited 2020 Apr 13]. Available online: <https://www.who.int/emergencies/diseases/novel-coronavirus-2019>
- Bi Q, Wu Y, Mei S, et al. Epidemiology and Transmission of COVID-19 in Shenzhen China: Analysis of 391 cases and 1,286 of their close contacts. *MedRxiv* 2020 [cited 2020 Apr 13]. Available online: <https://www.medrxiv.org/content/10.1101/2020.03.03.20028423v3>
- Wu Z, McGoogan JM. Characteristics of and important lessons from the coronavirus disease 2019 (COVID-19) outbreak in China: summary of a report of 72314 cases From the Chinese Center for Disease Control and Prevention. *JAMA* 2020;323:1239-42.
- Wang XF, Yuan J, Zheng Y, et al. Clinical and epidemiological characteristics of 34 children with 2019 novel coronavirus infection in Shenzhen. *Zhonghua Er Ke Za Zhi* 2020;58:E008.
- Chen F, Liu Z, Zhang F, et al. The first case of children with severe novel coronavirus pneumonia in China. *Chin J Pediatr* 2020;58.
- Lai CC, Shih TP, Ko WC, et al. Severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2) and corona virus disease-2019 (COVID-19): the epidemic and the challenges. *Int J Antimicrob Agents* 2020:105924.
- Han Q, Lin Q, Jin S, et al. Recent insights into 2019-nCoV: a brief but comprehensive review. *J Infect* 2020;S0163-4453(20)30087-6.
- Ferrara G, Zumla A, Maeurer M. Intravenous immunoglobulin (IVIg) for refractory and difficult-to-treat infections. *Am J Med* 2012;125:1036.e1-1036.e8.
- Björkander J, Nikoskelainen J, Leibl H, et al. Prospective open-label study of pharmacokinetics, efficacy and safety of a new 10% liquid intravenous immunoglobulin in patients with hypo-or agammaglobulinemia. *Vox Sanguinis* 2006;90:286-93.
- Crayne CB, Mitchell C, Beukelman T. Comparison of second-line therapy in IVIg-refractory Kawasaki disease: a systematic review. *Pediatr Rheumatol Online J* 2019;17:77.
- Hui DSC. An overview on severe acute respiratory syndrome (SARS). *Monaldi Arch Chest Dis* 2005;63(3).
- Zeng H. Diagnosis and treatment of severe acute respiratory syndrome in children. *J Appl Clin Pediatr* 2003;12:940-3.
- Guan W, Ni Z, Hu Y, et al. Clinical characteristics of 2019 novel coronavirus infection in China. *MedRxiv* 2020 [cited 2020 Apr 13]. Available online: <https://www.medrxiv.org/content/10.1101/2020.02.06.20020974v1>
- National Health Commission of the People's Republic of China. Diagnosis and treatments of pneumonia infected by novel coronavirus (trial version 7). 2020. [cited 2020 Mar 26]. Available online: <http://www.nhc.gov.cn/zyygj/s7653p>

- /202003/46c9294a7dfe4cef80dc7f5912eb1989.shtml
18. Ge L, Tian J, Li Y, et al. Association between prospective registration and overall reporting and methodological quality of systematic reviews: a meta epidemiological study. *J Clin Epidemiol* 2018;93:45-55.
  19. Moher D, Liberati A, Tetzlaff J, et al. PRISMA Group. Preferred Reporting Items for Systematic Reviews and Meta-analyses: the PRISMA statement. *BMJ* 2009;339:b2535.
  20. Higgins JPT, Thomas J, Chandler J, et al. *Cochrane Handbook for Systematic Reviews of Interventions* version 6.0 (updated July 2019). Cochrane, 2019. Available online: [www.training.cochrane.org/handbook](http://www.training.cochrane.org/handbook)
  21. National Institute for Health and Care Excellence 2003 [cited 2020 Mar 26]. Available online: <https://www.nice.org.uk/guidance/cg3/documents/appendix-4-quality-of-case-series-form2>
  22. The Joanna Briggs Institute. *The Joanna Briggs Institute Reviewers' Manual* 2016. Adelaide, SA Australia: The Joanna Briggs Institute, 2016.
  23. Stang A. Critical evaluation of the Newcastle-Ottawa scale for the assessment of the quality of nonrandomized studies in meta analyses. *Eur J Epidemiol* 2010;25:603-5.
  24. Nieva VF, Sorra J. Safety Culture Assessment: A Tool for Improving Patient Safety in Healthcare Organizations. *Quality and Safety in Health Care* 2003;12:ii17-23.
  25. Norris SL, Meerpohl JJ, Akl EA, et al. The skills and experience of GRADE methodologists can be assessed with a simple tool. *J Clin Epidemiol* 2016;79:150-8.e1.
  26. Guyatt GH, Oxman AD, Vist GE, et al. GRADE: an emerging consensus on rating quality of evidence and strength of recommendations. *BMJ* 2008;336:924-6.
  27. Guyatt GH, Oxman AD, Montori V, et al. GRADE guidelines: 5. Rating the quality of evidence—publication bias. *J Clin Epidemiol* 2011;64:1277-82.
  28. Guyatt GH, Oxman AD, Kunz R, et al. GRADE guidelines 6. Rating the quality of evidence—imprecision. *J Clin Epidemiol* 2011;64:1283-93.
  29. Guyatt GH, Oxman AD, Kunz R, et al. GRADE guidelines: 7. Rating the quality of evidence— inconsistency. *J Clin Epidemiol* 2011;64:1294-302.
  30. Guyatt GH, Oxman AD, Kunz R, et al. GRADE guidelines: 8. Rating the quality of evidence—indirectness. *J Clin Epidemiol* 2011;64:1303-10.
  31. Liu Y, Sun W, Li J, et al. Clinical features and progression of acute respiratory distress syndrome in coronavirus disease 2019. *MedRxiv* 2020 [cited 2020 Apr 13]. Available online: <https://www.medrxiv.org/content/10.1101/2020.02.17.20024166v3>
  32. Cao W, Liu X, Bai T, et al. High-dose intravenous immunoglobulin as a therapeutic option for deteriorating patients with Coronavirus Disease 2019. *Open Forum Infect Dis* 2020;7:ofaa102.
  33. Zeng H, Liu L, Wang B, et al. Clinical analysis on treatment of severe acute respiratory syndrome in children with intravenous Immunoglobulin. *J Clin Pediatr* 2003;21:533-4, 585.
  34. Wang JT, Sheng W, Fang C, et al. Clinical Manifestations, Laboratory Findings, and Treatment Outcomes of SARS Patients. *Emerg Infect Dis* 2004;10:818-24.
  35. Wu W, Wang J, Liu P, et al. Clinical features of 96 patients with severe acute respiratory syndrome from a hospital outbreak. *Zhonghua Nei Ke Za Zhi* 2003;42:453-7.
  36. Wu L, Xiao Z, Li Y. Intravenous Immunoglobulin was assisted treatment of 23 cases of severe infectious atypical pneumonia. *J Jinan Univ (Natural Science & Medicine Edition)* 2005;26:826-8.
  37. Stockman LJ, Bellamy R, Garner P. SARS: Systematic Review of Treatment Effects. *PLoS Med* 2006;3:e343.
  38. Sanders SL, Agwan S, Hassan M, et al. Immunoglobulin treatment for hospitalised infants and young children with respiratory syncytial virus infection. *Cochrane Database of Systematic Reviews* 2019;8:CD009417.
  39. Levy MM, Baylor MS, Bernard GR, et al. Clinical Issues and Research in Respiratory Failure from Severe Acute Respiratory Syndrome[J]. *Am J Respir Crit Care Med* 2005;171:518-26.
  40. Booth CM, Stewart TE. Severe acute respiratory syndrome and critical care medicine: The Toronto experience. *Crit Care Med* 2005;33:S53-60.
  41. Ohlsson A, Lacy JB. Intravenous immunoglobulin for suspected or proven infection in neonates. *Cochrane Database of Systematic Reviews* 2015;3:CD001239.
  42. Desborough MJ, Miller J, Thorpe SJ, et al. Intravenous immunoglobulin-induced haemolysis: a case report and review of the literature. *Transfus Med* 2014;24:219-26.
  43. Kazatchkine MD, Kaveri SV. Immunomodulation of autoimmune and inflammatory diseases with intravenous immune globulin. *N Engl J Med* 2001;345:747-55.
  44. Mustafa S, Balkhy H, Gabere MN. Current treatment options and the role of peptides as potential therapeutic components for Middle East Respiratory Syndrome (MERS): A review. *J Infect Public Health* 2018;11:9-17.
  45. Cabral D. A Guide to Contraindications to Childhood Vaccinations. *Can J Infect Dis* 2000;11:13-4.
  46. Guo Y, Tian X, Wang X, et al. Adverse Effects of

- Immunoglobulin Therapy. *Front Immunol* 2018;9:1299.
47. Chen Z, Fu J, Shu Q, et al. Diagnosis and treatment recommendation for pediatric coronavirus disease-19. *Zhejiang Da Xue Xue Bao Yi Xue Ban* 2020;49:1-8.
  48. Expert consensus on diagnosis and treatment of new coronavirus pneumonia in paediatrics of guangdong province. *Guangdong Med* 2020;41:217-21.
  49. Chinese society of pediatrics. Recommendations for diagnosis and prevention of 2019 novel coronavirus infection in children (1 edition). *Chin J Pediatr* 2020;58:169-74.
  50. The Efficacy of Intravenous Immunoglobulin Therapy for Severe 2019-nCoV Infected Pneumonia[Internet]. c2020 [cited 2020 Apr 13]. Available online: Available online: <https://clinicaltrials.gov/ct2/show/NCT04261426?cond=2019-nCoV&draw=2>

**Cite this article as:** Zhang J, Yang Y, Yang N, Ma Y, Zhou Q, Li W, Wang X, Huang L, Luo X, Fukuoka T, Ahn HS, Lee MS, Luo Z, Chen Y, Liu E, Yang K, Fu Z, on behalf of COVID-19 Evidence and Recommendations Working Group. Effectiveness of intravenous immunoglobulin for children with severe COVID-19: a rapid review. *Ann Transl Med* 2020;8(10):625. doi: 10.21037/atm-20-3305

**Search strategy****EMBASE**

- #1 'middle east respiratory syndrome coronavirus'/exp
- #2 'severe acute respiratory syndrome'/exp
- #3 'sars coronavirus'/exp
- #4 'COVID-19':ab,ti
- #5 'SARS-COV-2':ab,ti
- #6 'novel coronavirus':ab,ti
- #7 '2019-novel coronavirus':ab,ti
- #8 'coronavirus disease-19':ab,ti
- #9 'coronavirus disease 2019':ab,ti
- #10 'COVID 19':ab,ti
- #11 'novel cov':ab,ti
- #12 '2019-ncov':ab,ti
- #13 '2019-cov':ab,ti
- #14 'middle east respiratory syndrome':ab,ti
- #15 'middle east respiratory syndrome coronavirus':ab,ti
- #16 'mers':ab,ti
- #17 'mers-cov':ab,ti
- #18 'severe acute respiratory syndrome':ab,ti
- #19 'sars':ab,ti
- #20 'sars-cov':ab,ti
- #21 'sars-related':ab,ti
- #22 'sars-associated':ab,ti
- #23 #1-#22/ OR
- #24 'Immunoglobulins'/exp
- #25 'Intravenous Immunoglobulin\*':ab,ti
- #26 'Intravenous IG':ab,ti
- #27 'immune globulin\*':ab,ti
- #28 'IVIG':ab,ti
- #29 'IV Immunoglobulin\*':ab,ti
- #30 'Intravenous Antibodies':ab,ti
- #31 'gamma globulin\*':ab,ti
- #32 'gamma-globulin\*':ab,ti
- #33 'Flebogamma DIF':ab,ti
- #34 'Gamunex':ab,ti
- #35 'Globulin-N':ab,ti
- #36 'Globulin N':ab,ti
- #37 'Intraglobin':ab,ti
- #38 'Gammagard':ab,ti
- #39 'Gamimune':ab,ti
- #40 'Gamimmune':ab,ti
- #41 'Privigen':ab,ti
- #42 'Sandoglobulin':ab,ti
- #43 'Venoglobulin':ab,ti
- #44 'Iveegam':ab,ti

- #45 'Endobulin':ab,ti
- #46 'Gammonativ':ab,ti
- #47 #24-#46/OR
- #48 #23 AND #47

**PubMed**

- #1 "COVID-19" [Supplementary Concept]
- #2 "Severe Acute Respiratory Syndrome Coronavirus 2" [Supplementary Concept]
- #3 "Middle East Respiratory Syndrome Coronavirus" [Mesh]
- #4 "Severe Acute Respiratory Syndrome" [Mesh]
- #5 "SARS Virus" [Mesh]
- #6 "COVID-19" [Title/Abstract]
- #7 "SARS-COV-2" [Title/Abstract]
- #8 "Novel coronavirus" [Title/Abstract]
- #9 "2019-novel coronavirus" [Title/Abstract]
- #10 "coronavirus disease-19" [Title/Abstract]
- #11 "coronavirus disease 2019" [Title/Abstract]
- #12 "COVID 19" [Title/Abstract]
- #13 "Novel CoV" [Title/Abstract]
- #14 "2019-nCoV" [Title/Abstract]
- #15 "2019-CoV" [Title/Abstract]
- #16 "Middle East Respiratory Syndrome" [Title/Abstract]
- #17 "MERS" [Title/Abstract]
- #18 "MERS-CoV" [Title/Abstract]
- #19 "Severe Acute Respiratory Syndrome" [Title/Abstract]
- #20 "SARS" [Title/Abstract]
- #21 "SARS-CoV" [Title/Abstract]
- #22 "SARS-Related" [Title/Abstract]
- #23 "SARS-Associated" [Title/Abstract]
- #24 #1-#23/ OR
- #25 "Immunoglobulins, Intravenous" [Mesh]
- #26 "gamma-Globulins" [Mesh]
- #27 "Intravenous Immunoglobulin\*" [Title/Abstract]
- #28 "Intravenous IG" [Title/Abstract]
- #29 "immune globulin\*" [Title/Abstract]
- #30 IVIG [Title/Abstract]
- #31 "IV Immunoglobulin\*" [Title/Abstract]
- #32 "Intravenous Antibodies" [Title/Abstract]
- #33 "gamma globulin\*" [Title/Abstract]
- #34 "gamma-globulin\*" [Title/Abstract]
- #35 "Flebogamma DIF" [Title/Abstract]
- #36 Gamunex [Title/Abstract]
- #37 "Globulin-N" [Title/Abstract]
- #38 "Globulin N" [Title/Abstract]
- #39 Intraglobin [Title/Abstract]
- #40 Gammagard [Title/Abstract]
- #41 Gamimune [Title/Abstract]

- #42 Gamimmune [Title/Abstract]
- #43 Privigen [Title/Abstract]
- #44 Sandoglobulin [Title/Abstract]
- #45 Venoglobulin [Title/Abstract]
- #46 Iveegam [Title/Abstract]
- #47 Endobulin [Title/Abstract]
- #48 Gammonativ [Title/Abstract]
- #49 #25-#48/ OR
- #50 #24 AND #49

***Cochrane library***

- #1 MeSH descriptor: [Middle East Respiratory Syndrome Coronavirus] explode all trees
- #2 MeSH descriptor: [Severe Acute Respiratory Syndrome] explode all trees
- #3 MeSH descriptor: [SARS Virus] explode all trees
- #4 "COVID-19":ti,ab,ku=
- #5 "SARS-COV-2":ti,ab,kw
- #6 "Novel coronavirus":ti,ab,kw
- #7 "2019-novel coronavirus":ti,ab,kw
- #8 "Novel CoV":ti,ab,kw
- #9 "2019-nCoV":ti,ab,kw
- #10 "2019-CoV":ti,ab,kw
- #11 "coronavirus disease-19":ti,ab,kw
- #12 "coronavirus disease 2019":ti,ab,kw
- #13 "COVID 19":ti,ab,kw
- #14 "Middle East Respiratory Syndrome":ti,ab,kw
- #15 "MERS":ti,ab,kw
- #16 "MERS-CoV":ti,ab,kw
- #17 "Severe Acute Respiratory Syndrome":ti,ab,kw
- #18 "SARS":ti,ab,kw
- #19 "SARS-CoV":ti,ab,kw
- #20 "SARS-Related":ti,ab,kw
- #21 "SARS-Associated":ti,ab,kw
- #22 #1-#21/ OR
- #23 MeSH descriptor: [Immunoglobulins, Intravenous] explode all trees
- #24 MeSH descriptor: [gamma-Globulins] explode all trees
- #25 "Intravenous Immunoglobulin\*":ti,ab,kw
- #26 "Intravenous IG":ti,ab,kw
- #27 "immune globulin\*":ti,ab,kw
- #28 "IVIG":ti,ab,kw
- #29 "IV Immunoglobulin\*":ti,ab,kw
- #30 "Intravenous Antibodies":ti,ab,kw
- #31 "gamma globulin\*":ti,ab,kw
- #32 "gamma-globulin\*":ti,ab,kw
- #33 "Flebogamma DIF":ti,ab,kw
- #34 "Gamunex":ti,ab,kw
- #35 "Globulin-N":ti,ab,kw
- #36 "Globulin N":ti,ab,kw

- #37 "Intraglobin":ti,ab,kw
- #38 "Gammagard":ti,ab,kw
- #39 "Gamimmune":ti,ab,kw
- #40 "Gamimmune":ti,ab,kw
- #41 "Privigen":ti,ab,kw
- #42 "Sandoglobulin":ti,ab,kw
- #43 "Venoglobulin":ti,ab,kw
- #44 "Iveegam":ti,ab,kw
- #45 "Endobulin":ti,ab,kw
- #46 "Gammonativ":ti,ab,kw
- #47 #23-#46/ OR
- #48 #22 AND #47

### *Web of Science*

- #1 TOPIC: "COVID-19"
- #2 TOPIC: "SARS-COV-2"
- #3 TOPIC: "Novel coronavirus"
- #4 TOPIC: "2019-novel coronavirus"
- #5 TOPIC: "coronavirus disease-19" [Title/Abstract]
- #6 TOPIC: "coronavirus disease 2019" [Title/Abstract]
- #7 TOPIC: "COVID 19" [Title/Abstract]
- #8 TOPIC: "Novel CoV"
- #9 TOPIC: "2019-nCoV"
- #10 TOPIC: "2019-CoV"
- #11 TOPIC: " Middle East Respiratory Syndrome"
- #12 TOPIC: " MERS"
- #13 TOPIC: " MERS-CoV"
- #14 TOPIC: "Severe Acute Respiratory Syndrome"
- #15 TOPIC: "SARS"
- #16 TOPIC: "SARS-CoV"
- #17 TOPIC: "SARS-Related"
- #18 TOPIC: "SARS-Associated"
- #19 #1-#18 /OR
- #20 TOPIC: "Intravenous Immunoglobulin\*"
- #21 TOPIC: "Intravenous IG"
- #22 TOPIC: "immune globulin\*"
- #23 TOPIC: "IVIG"
- #24 TOPIC: "IV Immunoglobulin\*"
- #25 TOPIC: "Intravenous Antibodies"
- #26 TOPIC: "gamma globulin\*"
- #27 TOPIC: "gamma-globulin\*"
- #28 TOPIC: "Flebogamma DIF"
- #29 TOPIC: "Gamune"
- #30 TOPIC: "Globulin-N"
- #31 TOPIC: "Globulin N"
- #32 TOPIC: "Intraglobin"
- #33 TOPIC: "Gammagard"

- #34 TOPIC: "Gamimune"
- #35 TOPIC: "Gamimmune"
- #36 TOPIC: "Privigen"
- #37 TOPIC: "Sandoglobulin"
- #38 TOPIC: "Venoglobulin"
- #39 TOPIC: "Iveegam"
- #40 TOPIC: "Endobulin"
- #41 TOPIC: "Gammonativ"
- #42 #20-#41 /OR
- #43 #19 AND #42

### ***CBM***

- #1 " 新型冠状病毒 "[ 常用字段 : 智能 ]
- #2 "COVID-19"[ 常用字段 : 智能 ]
- #3 "COVID 19"[ 常用字段 : 智能 ]
- #4 "2019-nCoV"[ 常用字段 : 智能 ]
- #5 "2019-CoV"[ 常用字段 : 智能 ]
- #6 "SARS-CoV-2"[ 常用字段 : 智能 ]
- #7 " 中东呼吸综合征冠状病毒 "[ 不加权 : 扩展 ]
- #8 " 中东呼吸综合征 "[ 常用字段 : 智能 ]
- #9 "MERS"[ 常用字段 : 智能 ]
- #10 "MERS-CoV"[ 常用字段 : 智能 ]
- #11 " 严重急性呼吸综合征 "[ 不加权 : 扩展 ]
- #12 "SARS 病毒 "[ 不加权 : 扩展 ]
- #13 " 严重急性呼吸综合征 "[ 常用字段 : 智能 ]
- #14 "SARS"[ 常用字段 : 智能 ]
- #15 #1-#14/ OR
- #16 丙种球蛋白 [ 常用字段 : 智能 ]
- #17 静脉丙球 [ 常用字段 : 智能 ]
- #18 免疫球蛋白 [ 常用字段 : 智能 ]
- #19 IVIG[ 常用字段 : 智能 ]
- #20 " 免疫球蛋白类 "[ 不加权 : 扩展 ]
- #21 #16-#20/ OR
- #22 #15 AND #21

### ***Wangfang***

- #1 新型冠状病毒 [ 主题 ]
- #2 COVID-19[ 主题 ]
- #3 COVID 19[ 主题 ]
- #4 2019-nCoV[ 主题 ]
- #5 2019-CoV[ 主题 ]
- #6 SARS-CoV-2[ 主题 ]
- #7 中东呼吸综合征 [ 主题 ]
- #8 MERS[ 主题 ]
- #9 MERS-CoV[ 主题 ]
- #10 严重急性呼吸综合征 [ 主题 ]

- #11 SARS[ 主题 ]
- #12 #1-#11/ OR
- #13 丙种球蛋白 [ 主题 ]
- #14 静脉丙球 [ 主题 ]
- #15 IVIG[ 主题 ]
- #16 免疫球蛋白 [ 主题 ]
- #17 #13-#16/OR
- #18 #12 AND #17

### *CNKI*

- #1 " 新型冠状病毒 "[ 主题 ]
- #2 "COVID-19"[ 主题 ]
- #3 "COVID 19"[ 主题 ]
- #4 "2019-nCoV"[ 主题 ]
- #5 "2019-CoV"[ 主题 ]
- #6 "SARS-CoV-2"[ 主题 ]
- #7 " 中东呼吸综合征 "[ 主题 ]
- #8 "MERS"[ 主题 ]
- #9 "MERS-CoV"[ 主题 ]
- #10 " 严重急性呼吸综合征 "[ 主题 ]
- #11 "SARS"[ 主题 ]
- #12 #1-#11/ OR
- #13 " 丙种球蛋白 "[ 主题 ]
- #14 " 静脉丙球 "[ 主题 ]
- #15 " 免疫球蛋白 "[ 主题 ]
- #16 "IVIG"[ 主题 ]
- #17 #13-#16/ OR
- #18 #12 AND #17

**Table S1** The characteristics of excluded studies

No.	Title	Country	Journal	Year	Study type	Cause
1	Five cases of infant hematuria caused by human immunoglobulin	China	<i>Chinese Journal of Rural Medicine and Pharmacy</i>	2005	Case report	Intramuscular injection of IVIG
2	SARS: systematic review of treatment effects	US	<i>PLoS Medicine</i>	2006	Review	Review
3	A hospital outbreak of severe acute respiratory syndrome in Guangzhou, China	China	<i>Chinese medical journal</i>	2003	Case series	Without the outcomes of efficacy and safety of IVIG
4	Comparison of clinical course of patients with severe acute respiratory syndrome among the multiple generations of nosocomial transmission	China	<i>Chinese Medical Journal</i>	2004	Case series	Without the outcomes of efficacy and safety of IVIG
5	Evaluation of the efficacy and safety of corticosteroid in the treatment of severe SARS in Guangdong province with multi-factor regression analysis	China	<i>Chinese Critical Care Medicine</i>	2008	Regression analysis	Without the outcomes of efficacy and safety of IVIG
6	The therapeutic effect of high flow nasal cannula oxygen therapy for the first imported case of Middle East respiratory syndrome to China	China	<i>Chinese Critical Care Medicine</i>	2015	Case report	Without the outcomes of efficacy and safety of IVIG
7	Clinical analysis of pediatric SARS cases in Beijing	China	<i>Chinese Journal of Pediatrics</i>	2003	Case series	Without the outcomes of efficacy and safety of IVIG
8	Clinical analysis of the first patient with imported Middle East respiratory syndrome in China	China	<i>Chinese Critical Care Medicine</i>	2015	Case report	Without the outcomes of efficacy and safety of IVIG
9	Clinical characteristics and therapy of severe acute respiratory syndrome	China	<i>Chinese Famous Doctor Forum</i>	2004	Case series	Full-text unavailable
10	Multivariable analysis of factors affecting clinical course in patients with SARS	China	<i>Journal of Sun Yat-sen University (Medical Sciences)</i>	2004	Regression analysis	Without the outcomes of efficacy and safety of IVIG
11	Clinical analysis of 136 cases of severe acute respiratory syndrome	China	<i>Chinese Journal of Respiratory and Critical Care Medicine</i>	2003	Case series	Without the outcomes of efficacy and safety of IVIG
12	Evaluation of the efficacy and safety of corticosteroid in the treatment of severe SARS in Guangdong province with multi-factor regression analysis	China	<i>Chinese Critical Care Medicine</i>	2008	Regression analysis	Without the outcomes of efficacy and safety of IVIG
13	Study of the clinical diagnosis and treatment of the severe acute respiratory syndromes	China	<i>Jiangsu Medical Journal</i>	2003	Case report	Without the outcomes of efficacy and safety of IVIG
14	Can immunoglobulin, thymosin, and interferon protect against SARS?	China	<i>Hohhot Technology Journal</i>	2003	Science & Technology Daily	Review
15	The search for therapeutic options for Middle East respiratory syndrome (MERS)	Saudi Arabia	<i>Journal of Infection and Public Health</i>	2016	Editorial	Review
16	Clinical findings, treatment and prognosis in patients with severe acute respiratory syndrome (SARS)	China	<i>Journal of the Chinese Medical Association</i>	2005	Editorial	Review
17	Intravenous immunoglobulin G is remarkably beneficial in chronic immune dysschwannian/dysneuronal polyneuropathy, diabetes-2 neuropathy, and potentially in severe acute respiratory syndrome	United States	<i>Acta Myologica</i>	2003	Review	Full-text unavailable
18	Severe acute respiratory syndrome: public health response and clinical practice update for an emerging disease	United States	<i>Current Opinion in Pediatrics</i>	2004	Review	IVIG not mentioned
19	Treatment of severe acute respiratory syndrome	Hong Kong	<i>European Journal of Clinical Microbiology and Infectious Diseases</i>	2005	Review	IVIG not mentioned
20	Current treatment options and the role of peptides as potential therapeutic components for Middle East respiratory syndrome (MERS): a review	Saudi Arabia	<i>Journal of Infection and Public Health</i>	2018	Review	Review
21	Management of hospital-acquired severe acute respiratory syndrome with different disease spectrum	China	<i>Journal of the Chinese Medical Association: JCMA</i>	2003	Case report	Without the outcomes of efficacy and safety of IVIG
22	Neurological manifestations in severe acute respiratory syndrome	China	<i>Acta Neurologica Taiwanica</i>	2005	Review	Review
23	Diagnosis and treatment of severe acute respiratory syndrome in children	China	<i>Journal of Applied Clinical Pediatrics</i>	2003	Medical advice	Review
24	Experience of INF- $\alpha$ for treating severe acute respiratory distress syndrome	China	<i>Journal of Modern Medicine &amp; Health</i>	2004	Case series	Without the outcomes of efficacy and safety of IVIG
25	Study of severe acute respiratory syndrome in Shantou	China	<i>Journal of Shantou University Medical College</i>	2003	Case report	Without the outcomes of efficacy and safety of IVIG

**Table S2** GRADE evidence profile

No. of studies	Sample size	Certainty assessment					Effect value (95% CI)	Certainty
		Risk of bias	Inconsistency	Indirectness	Imprecision	Other considerations		
Serum globulin								
RCT (31)	44	Serious <sup>1</sup>	Not applicable	Serious <sup>2</sup>	Not serious	None	MD 6.00 (4.75 to 7.25)	⊕⊕○○ low
Nosocomial infection rate								
RCT (31)	44	Serious <sup>1</sup>	Not applicable	Serious <sup>2</sup>	Not serious	None	OR 1.25 (0.47 to 3.31)	⊕⊕○○ low
The risk of death								
RCT (31)	44	Serious <sup>1</sup>	Not applicable	Serious <sup>2</sup>	Not serious	None	OR 1.70 (0.51 to 0.73)	⊕⊕○○ low

Explanations: <sup>1</sup>, unclear risk of bias in allocation concealment, random sequence generation and blinding; <sup>2</sup>, using other drugs (such as interferon, hormone, etc.) before intervention. CI, confidence interval; OR, odds ratio; MD, mean deviation.