



Evaluation of the roles and regulatory mechanisms of PD-1 target molecules in NSCLC progression

Yun-Qiang Zhang^{1#}, Ye Yuan^{2#}, Jun Zhang², Cheng-Yi Lin², Jia-Long Guo², Hua-Song Liu², Qiang Guo^{2,3}

¹Department of Thoracic Surgery, Beilun District People's Hospital of Ningbo, Ningbo, China; ²Department of Cardiothoracic Surgery, Taihe Hospital, Hubei University of Medicine, Shiyan, China; ³Department of Thoracic Surgery, Union Hospital, Tongji Medical College, Huazhong University of Science and Technology, Wuhan, China

Contributions: (I) Conception and design: Q Guo, HS Liu; (II) Administrative support: JL Guo, Y Yuan; (III) Provision of study materials or patients: YQ Zhang; (IV) Collection and assembly of data: Y Yuan; (V) Data analysis and interpretation: J Zhang, CY Lin; (VI) Manuscript writing: All authors; (VII) Final approval of manuscript: All authors.

[#]These authors contributed equally to this work.

Correspondence to: Qiang Guo; Hua-Song Liu. Department of Cardiothoracic Surgery, Taihe Hospital, Hubei University of Medicine, Shiyan 442012, China. Email: guoqianglidan@163.com; huasong819@qq.com.

Background: Targeted programmed cell death protein 1 (PD-1) therapy could effectively improve the long-term prognosis of patients with non-small cell lung cancer (NSCLC). The role of PD-1 targets in the progression of NSCLC has not been fully revealed.

Methods: The differentially expressed genes (DEGs) in patients' blood after NSCLC treatment with PD-1 blocker nivolumab in the GSE141479 dataset were analyzed by GEO2R and identified in the TCGA database. The mechanism of action involved in the PD-1 target molecules via the Gene Ontology (GO) and Kyoto Encyclopedia of Genes and Genomes (KEGG). The protein-protein interaction (PPI) network shows the relationship between PD-1 target molecules. The factors affecting the prognosis of NSCLC patients were identified via the COX regression analysis and survival analysis to build the risk model and nomogram.

Results: There were 64 DEGs in patients' blood after nivolumab treatment and 48 DEGs in NSCLC tissues. The PD-1 target molecules involved cell proliferation, DNA replication, cell cycle, lung cancer, and other cellular processes. The prognostic factors CCNA2, CHEK1, DLGAP5, E2F8, FOXM1, HIST1H2BH, HJURP, MKI67, PLK1, TPX2, and TYMS, and the independent factors HIST1H2BH and PLK1, influenced the prognosis of NSCLC patients. HIST1H2BH and PLK1 were overexpressed in LUAD and LUSC tissues. The elevated expression levels of HIST1H2BH and PLK1 were related to the overall survival (OS) and the progression-free survival of NSCLC patients. High-risk NSCLC patients had a poor prognosis and were an independent factor influencing the poor prognosis of NSCLC patients. The high-risk model group was enriched with signaling mechanisms such as cell cycle, DNA replication, and homologous recombination.

Conclusions: The risk model based on PD-1 target molecules was helpful to assess the prognosis of NSCLC patients. HIST1H2BH and PLK1 might become prognostic biomarkers of NSCLC patients.

Keywords: Programmed cell death protein 1 (PD-1); non-small cell lung cancer (NSCLC); nivolumab

Submitted May 25, 2021. Accepted for publication Jul 08, 2021.

doi: 10.21037/atm-21-2963

View this article at: <https://dx.doi.org/10.21037/atm-21-2963>

Introduction

In recent years, programmed cell death protein 1 (PD-1) has been one of the inhibitory immune checkpoints that activate T cells, NK cells, and B cells. PD-1 interacts with programmed death ligand 1 (PD-L1) to produce inhibitory signaling transduction, resulting in anti-tumor effects during cancer treatment (1). For example, in MKN-45 and MGC-803 cells of gastric cancer (GC), anti-cytotoxic T lymphocyte antigen-4 (anti-CTLA-4) and anti-PD-1 combined therapy could significantly inhibit cell proliferation, migration, invasion, and epithelial-mesenchymal transition (EMT), as well as induce cell apoptosis *in vitro*, and inhibit tumor formation *in vivo*. Mechanism studies have shown that the anti-CTLA-4 and anti-PD-1 combined therapy could inhibit the activation of β -catenin, MAPK, and PI3K/AKT signaling pathways. Interference with PD-1 expression inhibited GC cell proliferation, migration, invasion, and EMT and induced cell apoptosis (2). Inhibition of BET protein expression inhibited the expression level of PD-1/PD-L1 in triple-negative breast cancer. BET protein regulated the PD-1 expression level, reduced the production and signaling transduction of interferon- γ in the T cells, and inhibited the expression of PD-L1 in breast cancer cells. Inhibition of BET protein improved tumor cell-specific T cell toxicity (3). In non-small cell lung cancer (NSCLC) tissues, PD-L1 expression was positive in 32.6% of patients. In patients with high-grade malignancy and lymph node metastasis, the expression level of PD-L1 increased significantly (4). In lung adenocarcinoma (LUAD), patients with low PD-L1 expression levels had a longer overall survival (OS) (4). Cyclin-dependent kinase 7 (CDK7) mRNA and protein expression levels were related to the poor prognosis of NSCLC patients. The decrease of CDK7 expression level could cause NSCLC cell apoptosis and inhibit tumor growth. The CDK7 inhibitor THZ1 could down-regulate PD-L1 expression level by inhibiting MYC activity. THZ1 enhanced anti-tumor immunity by recruiting CD8 T cell infiltration and synergistically with anti-PD-1 therapy (5). This indicated that the anti-PD-1 treatment of cancer patients had important value.

Nivolumab is the PD-1 inhibitor that binds to PD-L1 to the surface of T cells to block the immunosuppressive signaling pathway triggered by PDL-1/2 and restore the anti-tumor function of T cells. Currently, nivolumab is used to treatment the patients with unresectable or metastatic melanoma and advanced lung squamous cell carcinoma

(LUSC) and show the safe and effective therapeutic effect (6-9). For example, compared with the cytotoxic T-cell antigen-4 (CTLA-4) blocker ipilimumab, nivolumab has sustained recurrence-free survival benefits in patients with stage IIIB-C or IV melanoma after surgery. Nivolumab is also an effective adjuvant treatment for patients with high-risk melanoma after surgery (6). At present, many studies had shown that the PD-1 inhibitor nivolumab could improve the prognosis of cancer patients (8,10,11). For example, Fujimoto *et al.* reported that the 6-month PFS rate of nivolumab in patients with advanced NSCLC was 56%, the remission rate was 39%, and the disease control rate was 72%. There were no treatment-related deaths (8). Sugawara *et al.* reported that the median PFS of the nivolumab group was higher than that of the placebo group. Nivolumab treatment improved the PFS and objective remission rate of patients with advanced non-squamous NSCLC (10). Hoffner *et al.* reported that the combination of anti-PD-L1 immunotherapy and platinum-based chemotherapy can improve the prognosis of NSCLC and SCLC patients (11). Nivolumab has been effective therapeutic value in NSCLC treatment, and the mechanisms in anti-NSCLC treatment have not been fully revealed. Therefore, differentially expressed genes (DEGs) in the blood of patients from the GSE141479 dataset were analyzed after nivolumab treatment (12), the mechanisms of PD-1 target molecules in the progression of NSCLC were explored, and hub target molecules were screened in order to construct a risk model and nomogram to assess the prognosis of NSCLC patients. This could provide a theoretical reference point and new approach for cancer treatment.

We present the following article in accordance with the REMARK reporting checklist (available at <https://dx.doi.org/10.21037/atm-21-2963>).

Methods

The study was conducted in accordance with the Declaration of Helsinki (as revised in 2013).

GSE141479 dataset

The GSE141479 dataset of the Gene Expression Omnibus [GEO (<http://www.ncbi.nlm.nih.gov/geo/>)] database contained gene expression data from the blood samples of 74 NSCLC patients. Among them, 33 NSCLC patients were the blood samples treated with PD-1 inhibitor nivolumab. The platform information was GPL23126

[Clariom_D_Human] Affymetrix Human Clariom D Assay [transcript (gene) version]. The gene expression levels were explored in blood samples of patients with NSCLC after nivolumab treatment via the GEO2R. Screening criteria: $\log_{2}FC > 0.585$ or $\log_{2}FC < -0.585$ and $P < 0.05$.

The Cancer Genome Atlas (TCGA) database

The gene expression data of 108 normal lung tissue samples and 1,037 NSCLC tissue samples, the prognosis and clinicopathological characteristics data of NSCLC patients were downloaded from the TCGA (<https://portal.gdc.cancer.gov/projects/>) website. The limma package was used to analyze the expression levels of PD-1 target molecules in normal lung tissue and cancer tissue ($\log_{2}FC > 1$ or $\log_{2}FC < -1$ and $P < 0.05$).

DAVID database

The DAVID database could analyze biological processes (BP), molecular functions (MF), cellular components (CC), signaling mechanisms, and the transcription factor regulation involved in multiple genes. In the DAVID database, the gene ontology (GO), Kyoto encyclopedia of genes and genomes (KEGG), and transcription factor analysis were used to analyze the MF, BP, CC, and signaling mechanisms involved in the abnormally expressed PD-1 target molecules of the TCGA database.

Protein-protein interaction (PPI) network

The abnormally expressed PD-1 target molecules in the TCGA database were entered into the Search Tool for the Retrieval of Interacting Genes (STRING) database to explore the PPI. The combination score > 0.4 was the screening criterion. Cytoscape 3.6.1 software was used to visualize the relationship between proteins in the PPI network, and the MCODE method was used for enrichment analysis.

The risk model and nomogram

The prognostic data of NSCLC in the TCGA database were sorted out, and missing values were deleted. PD-1 target gene expression and prognosis of NSCLC patients was spliced. Univariate Cox regression analysis evaluated the impact of PD-1 target molecules on the prognosis of NSCLC patients. Multivariate Cox regression analysis

and the AIC method were used to screen independent factors affecting the prognosis of NSCLC patients, and a risk model was constructed. Kaplan-Meier (K-M) survival analysis and nomogram showed the adverse prognosis of NSCLC patients in the high- and low-risk groups.

UALCAN database

The expression levels of HIST1H2BH and PLK1 were explored in NSCLC subtypes LUAD and LUSC tissues by using the UALCAN database. The relationship between the expression levels of HIST1H2BH and PLK1 and the clinicopathological characteristics of LUAD and LUSC patients were explored via correlation analysis. Screening criteria: $P < 0.05$.

K-M Plotter database

K-M Plotter database was a commonly used prognostic analysis database. This database contained prognostic information for patients with lung cancer, breast cancer, and stomach cancer. In the K-M Plotter database, the median value of HIST1H2BH and PLK1 expression levels were grouped to evaluate their potential role in the prognosis of NSCLC patients.

CbioPortal database

The CbioPortal database provided large-scale cancer genomics datasets and visual analysis functions. In the CbioPortal database, the TCGA NSCLC data was used to show the mutations of HIST1H2BH and PLK1 in NSCLC and its subtypes LUAD and LUSC.

Lung cancer explore (LCE) database

The LCE database covered lung cancer gene expression data and prognostic information, and meta-analysis could be used to visualize gene expression in cancer patient tissues and its prognostic value. In the LCE database, the expression and prognostic value of HIST1H2BH and PLK1 were explored in lung cancer and its subtypes LUAD and LUSC via the meta-analysis.

GEPIA database

The data in the GEPIA database came from the TCGA and GTEx databases. Correlation analysis was used to explore

the relationship between the expression level of PD-1 and the expression levels of risk model factors in normal lung tissues, LUAD, and LUSC tissues.

The signaling mechanisms of the risk model

Gene Set Enrichment Analysis (GSEA) was commonly used to evaluate the signaling mechanisms involved in influencing factors. In this study, we divided the TCGA NSCLC gene expression data into high- and low-risk groups based on the median value of risk scores, and the signaling mechanisms involved in the high- and low-risk groups were explored via the GSEA (13). The program was run for 1,000 cycles, and nominal P value (NOM P) <0.05 was used as the screening criterion.

Statistics analysis

Blood of patients with NSCLC after nivolumab treatment was analyzed for DEGs using the GEO2R. The expression levels of PD-1 target molecules in NSCLC tissues were analyzed by the R limma package. Grouping criteria: median value of gene expression and risk score. Univariate and multivariate Cox regression analysis explored potential factors affecting the prognosis of NSCLC patients, including age, gender, clinical stage, T stage, lymph node metastasis, distant metastasis, high- and low-risk. K-M survival analysis showed the survival curve of NSCLC patients with high and low risk. P<0.05 was considered statistically significant.

Results

The expression levels of PD-1 target molecules in NSCLC

Compared with 41 NSCLC patients, there were 64 DEGs in the blood of 33 NSCLC patients who received PD-1 inhibitor treatment (P<0.05). There are 11 highly expressed genes and 53 lowly expressed genes (Figure 1 and Table 1). In the TCGA database, compared with normal lung tissues, the expression levels of 48 PD-1 target molecules in NSCLC tissues were significantly increased (Figure 2 and Table 2).

The biological functions and signaling mechanisms of PD-1 target molecules

GO analysis showed that the PD-1 target molecules were involved in the mitotic nuclear division, chromosome

segregation, cell division, cell proliferation, DNA binding, protein kinase binding, DNA replication, mitotic cytokinesis, DNA repair, and other cellular functions (Figure 3A,B,C and Table S1). KEGG signaling mechanism analysis showed that the PD-1 target molecules were involved in viral carcinogenesis, systemic lupus erythematosus, alcoholism, cell cycle, and progesterone-mediated oocyte maturation (Figure 3D). In addition, PD-1 target molecules were also involved in breast cancer, ovarian cancer, leukemia, bladder cancer, chronic obstructive pulmonary disease, esophageal adenocarcinoma, lung cancer, head and neck cancer, and other diseases (Figure S1 and Table 3).

The network of PD-1 target molecules

In the DAVID database, there were 3 transcription factors (NFY, E2F, and MEF2) enriched by 48 PD-1 target molecules (P<0.05). Figure S2 showed the relationship between transcription factors NFY, E2F, MEF2, and PD-1 target molecules, and the PPI network was used to visualize the relationship between the proteins of 48 PD-1 target molecules (Figure 4).

Construction of PD-1 target molecular risk model

Univariate Cox regression analysis showed that CCNA2, CHEK1, DLGAP5, E2F8, FOXM1, HIST1H2BH, HJURP, MKI67, PLK1, TPX2, and TYMS were the prognostic factors of NSCLC patients (Figure 5A and Table 4). On this basis, multivariate Cox regression analysis and AIC screening were performed, and it was found that HIST1H2BH and PLK1 were independent factors influencing the prognosis of NSCLC patients (Figure 5B and Table 5). Figure 5C and 5D showed the high and low scores and survival status of NSCLC patients.

The value of risk model factors was verified in NSCLC

In the UALCAN database, HIST1H2BH was overexpressed in NSCLC subtype LUAD tissues. The expression level of HIST1H2BH was correlated with the age, gender, smoking history, and histological subtype of LUAD patients (Figure 6A and Table 6). PLK1 was overexpressed in LUAD tissues. The expression level of PLK1 was correlated with the age, gender, clinical stage, smoking history, histological subtype, and TP53 mutation status of LUAD patients (Figure 6B and Table 6). HIST1H2BH was overexpressed in NSCLC subtype LUSC tissues. The expression level




Figure 1 The expression level of the PD-1 target molecules in GSE141479 NSCLC blood samples. (A) Box plot showing that the samples have good consistency; (B) the expression density of sample genes; (C) highly expressed genes; (D) lowly expressed genes. PD-1, programmed cell death protein 1; NSCLC, non-small cell lung cancer.

of HIST1H2BH was correlated with smoking history and histological subtypes in LUSC patients (*Table 7*). PLK1 was overexpressed in LUSC tissues. The expression level of PLK1 was related to the age, gender, clinical stage, smoking history, lymph node metastasis, and TP53 mutation status of LUSC patients (*Table 7*). In the K-M Plotter database, K-M survival analysis showed that elevated expression levels of HIST1H2BH and PLK1 were related to the OS and PFS of NSCLC patients (*Figure 7*).

In addition, the meta-analysis results showed that HIST1H2B and PLK1 were overexpressed in lung cancer tissues, and the hazard ratio (HR) were 1.18 and 1.71, respectively, and they were statistically significant (*Figure 8A,B*). The overexpression of HIST1H2BH and PLK1 indicated that the prognosis of lung cancer patients was not

better. The HR was 1.05 and 1.16. There was no statistical significance between the overexpression of HIST1H2BH and the prognosis of lung cancer patients (*Figure 8C,D*). The results of subgroup analysis showed that HIST1H2BH was overexpressed in LUAD and LUSC tissues, with HR of 1.27 and 2.1, respectively, which were statistically significant (*Figure S3A*). PLK1 was overexpressed in LUAD and LUSC tissues, and the HR was 1.79 and 3.15, respectively, which were statistically significant (*Figure S3B*). The overexpression of HIST1H2BH suggested that the prognosis of LUAD patients was poor, and the HR was 1.08, which is statistically significant (*Figure S4A*). The low expression of HIST1H2BH indicated a poor prognosis for LUSC patients, and the HR was 0.89, which was statistically significant (*Figure S4A*). The overexpression of PLK1

Table 1 The expression level of PD-1 target molecules in GSE141479 NSCLC blood samples

Gene	P	logFC
<i>SNORD111</i>	0.00493817	0.72559394
<i>RNU6-948P</i>	0.03444443	0.69960397
<i>KCNQ1OT1_5</i>	0.00166819	0.68030422
<i>LEF1-AS1</i>	0.03273117	0.65527865
<i>SNORD63</i>	0.00920666	0.64174079
<i>GPRASP1</i>	0.00110561	0.63013588
<i>HMGB3P4</i>	0.00121729	0.61194145
<i>MTCO1P31</i>	0.02050908	0.60380088
<i>ARHGAP17</i>	0.0023222	0.603064
<i>CUBN</i>	0.01017528	0.59152504
<i>TRAJ14</i>	0.002776	0.58637359
<i>CDCA2</i>	0.01072148	-0.59144421
<i>FAM83D</i>	0.00157576	-0.59256397
<i>FOXM1</i>	0.01552184	-0.59382417
<i>H2BFS</i>	0.00312788	-0.6007239
<i>KIF11</i>	0.01282965	-0.60308543
<i>CENPU</i>	0.00655007	-0.6075029
<i>MLC1</i>	0.03826616	-0.60756707
<i>NUSAP1</i>	0.00545245	-0.61177406
<i>SGOL1</i>	0.00965143	-0.61373539
<i>MCM2</i>	0.00100469	-0.61971768
<i>CDC6</i>	0.02287365	-0.62300489
<i>KIF20A</i>	0.01558623	-0.62524289
<i>GMNN</i>	0.00932813	-0.6271711
<i>CENPM</i>	0.00565937	-0.6351112
<i>RFC2</i>	0.00022004	-0.63632544
<i>CENPN</i>	0.00127519	-0.63697021
<i>BRCA2</i>	0.00532567	-0.64152522
<i>E2F8</i>	0.03568498	-0.64326644
<i>EXO1</i>	0.00176873	-0.65220136
<i>HIST1H2BI</i>	0.00253962	-0.65270249
<i>HIST1H2BJ</i>	0.00212488	-0.65351784
<i>PLK1</i>	0.02763867	-0.65474106
<i>DTL</i>	0.03959844	-0.65616881

Table 1 (continued)**Table 1** (continued)

Gene	P	logFC
<i>HIST1H2BK</i>	0.00185741	-0.65692162
<i>CENPF</i>	0.01276753	-0.67133204
<i>CDKN3</i>	0.03077191	-0.6757393
<i>HIST1H2BH</i>	0.0023265	-0.67885224
<i>CKS2</i>	0.00375082	-0.69673774
<i>PRRG4</i>	0.00719681	-0.70325329
<i>HJURP</i>	0.00669842	-0.70537786
<i>STMN1</i>	0.00217296	-0.70766827
<i>PRC1</i>	0.00168319	-0.7179418
<i>CCNB2</i>	0.03641449	-0.71887783
<i>NDC80</i>	0.00073926	-0.72614622
<i>TPX2</i>	0.01299743	-0.74730387
<i>KIF15</i>	0.00218344	-0.74827382
<i>HIST1H2BL</i>	0.00364301	-0.7494745
<i>RAD51AP1</i>	0.00063054	-0.7701625
<i>HIST1H3B</i>	0.00397439	-0.78280432
<i>CCNA2</i>	0.02322892	-0.79709647
<i>CD38</i>	0.01137765	-0.79908712
<i>HIST1H2BB</i>	0.00085693	-0.80144284
<i>HIST1H1B</i>	0.00378843	-0.80473676
<i>CENPW</i>	0.00594784	-0.80853294
<i>TYMS</i>	0.01973621	-0.81167961
<i>TOP2A</i>	0.00627913	-0.84272863
<i>HIST1H2BM</i>	0.01649506	-0.88164367
<i>DLGAP5</i>	0.03435208	-0.90377947
<i>CHEK1</i>	0.00301191	-0.92741563
<i>MELK</i>	0.0003221	-0.93779218
<i>FAM111B</i>	0.00026547	-0.97286209
<i>MKI67</i>	0.00484638	-0.99910995
<i>KIAA0101</i>	0.00937747	-1.04596747

PD-1, programmed cell death protein 1; NSCLC, non-small cell lung cancer.




Figure 2 The expression level of PD-1 target molecules in the TCGA NSCLC tissues. (A) Heat map representing the DEGs; (B) violin chart demonstrating the top 10 DEGs via the fold-change. *******, $P < 0.001$. PD-1, programmed cell death protein 1; TCGA, the cancer genome atlas; NSCLC, non-small cell lung cancer; DEGs, differentially expressed genes.

Table 2 The expression levels of PD-1 target molecules in NSCLC tissues

Gene	logFC	P
<i>HIST1H3B</i>	6.971563358	5.22E-42
<i>HIST1H1B</i>	6.889830378	1.98E-34
<i>HIST1H2BB</i>	6.6176645	2.92E-16
<i>HIST1H2BM</i>	6.546656387	1.70E-27
<i>HIST1H2BI</i>	6.144328211	3.81E-19
<i>HIST1H2BH</i>	5.25752465	1.97E-39
<i>HJURP</i>	4.423193962	9.76E-63
<i>TPX2</i>	4.356893524	1.49E-61
<i>MELK</i>	4.28293495	2.60E-61
<i>TOP2A</i>	4.239513136	6.71E-63
<i>FOXM1</i>	4.206146682	1.40E-60
<i>EXO1</i>	4.176032779	4.79E-62
<i>DLGAP5</i>	4.123032333	1.54E-61
<i>CENPF</i>	4.004101702	2.23E-61
<i>CDC6</i>	3.962775113	8.09E-62
<i>PLK1</i>	3.804871556	1.14E-62
<i>CCNB2</i>	3.791913823	1.42E-61
<i>HIST1H2BL</i>	3.771763049	4.85E-33
<i>NDC80</i>	3.66052765	1.20E-60
<i>CDCA2</i>	3.618670951	2.97E-60
<i>KIF20A</i>	3.560206265	2.46E-61
<i>CCNA2</i>	3.560063035	9.08E-61
<i>MKI67</i>	3.512151918	2.23E-58
<i>CDKN3</i>	3.434927857	1.04E-57

Table 2 (continued)

Table 2 (continued)

Gene	logFC	P
<i>FAM83D</i>	3.412001161	2.99E-50
<i>E2F8</i>	3.334165686	2.05E-61
<i>PRC1</i>	3.322365587	1.58E-61
<i>KIF11</i>	3.262694147	7.05E-62
<i>NUSAP1</i>	3.233047006	1.10E-60
<i>KIF15</i>	3.110763871	7.51E-59
<i>FAM111B</i>	3.110199706	4.99E-57
<i>MCM2</i>	2.988886147	5.17E-58
<i>CENPU</i>	2.922017757	3.96E-59
<i>RAD51AP1</i>	2.910137622	4.25E-55
<i>CHEK1</i>	2.741114388	2.08E-61
<i>HIST1H2BJ</i>	2.624552264	7.72E-28
<i>CENPM</i>	2.594186936	9.10E-51
<i>CENPW</i>	2.583310716	1.63E-42
<i>DTL</i>	2.466385804	3.72E-50
<i>TYMS</i>	2.401391345	4.02E-56
<i>GMNN</i>	1.900079663	1.15E-56
<i>CENPN</i>	1.899039582	1.93E-47
<i>STMN1</i>	1.625279048	2.36E-35
<i>CKS2</i>	1.53273206	6.79E-34
<i>BRCA2</i>	1.461967704	6.31E-37
<i>HIST1H2BK</i>	1.422823205	4.16E-26
<i>RFC2</i>	1.239427886	1.96E-48
<i>H2BFS</i>	1.214789768	0.02134121

PD-1, programmed cell death protein 1; NSCLC, non-small cell lung cancer.




Figure 3 The biological functions and signaling mechanisms of PD-1 target molecules via the GO and KEGG. (A) BP; (B) CC; (C) MF; (D) KEGG. GO, gene ontology; BP, biological processes; CC, cellular component; MF, molecular function; KEGG, Kyoto encyclopedia of genes and genomes; PD-1, programmed cell death protein 1.

Table 3 Disease related to PD-1 target molecules

Category	Term	Count	P
GAD_DISEASE	Breast Cancer	10	2.72E-05
GAD_DISEASE	Longevity	9	1.68E-05
GAD_DISEASE	Chronic renal failure Kidney Failure, Chronic	8	0.011029875
GAD_DISEASE	Ovarian cancer	7	4.88E-04
GAD_DISEASE	Breast cancer	7	0.002764474
GAD_DISEASE	Bladder Cancer	7	0.003069375
GAD_DISEASE	Chronic obstructive pulmonary disease	6	0.005391657
GAD_DISEASE	Lung cancer	6	0.013295264
GAD_DISEASE	Lung Cancer	6	0.022918521
GAD_DISEASE	Leukemia, Lymphocytic, Chronic, B-Cell	5	0.002712277
GAD_DISEASE	Esophageal adenocarcinoma	5	0.015692824
GAD_DISEASE	Pharmacogenetic studies	4	0.003197262
GAD_DISEASE	Adenocarcinoma Pancreatic Neoplasms	3	0.009443208
GAD_DISEASE	Head and neck cancer	3	0.026277742
GAD_DISEASE	Abortion, Spontaneous	3	0.036618707

PD-1, programmed cell death protein 1.




Figure 4 The PPI network of 48 PD-1 target molecules. (A) PPI network; (B,C,D) the enrichment analysis by MCODE. PPI, protein-protein interaction; PD-1, programmed cell death protein 1.




Figure 5 The risk model of PD-1 target molecules. (A) Univariate Cox regression analysis of the prognostic factors; (B) heat map showing the relationship between risk and prognostic factors; (C,D) the risk scores and survival status of NSCLC patients. PD-1, programmed cell death protein 1; NSCLC, non-small cell lung cancer.

Table 4 Univariate Cox regression analysis showed the prognostic factors of NSCLC patients

Gene	HR	HR.95L	HR.95H	P
<i>CCNA2</i>	1.119030214	1.015250316	1.233418597	0.023527732
<i>CHEK1</i>	1.161268534	1.003467151	1.343885155	0.044812367
<i>DLGAP5</i>	1.120187116	1.007468832	1.24551662	0.035951079
<i>E2F8</i>	1.188481809	1.015445446	1.391004328	0.031485368
<i>FOXM1</i>	1.105655354	1.017333811	1.20164468	0.018052236
<i>HIST1H2BH</i>	0.896796742	0.827448404	0.971957155	0.007986484
<i>HJURP</i>	1.113339724	1.001826182	1.237265867	0.046168795
<i>MKI67</i>	1.118363726	1.008411159	1.24030502	0.034124878
<i>PLK1</i>	1.184761851	1.061018977	1.322936416	0.002592543
<i>TPX2</i>	1.102047152	1.014411384	1.19725384	0.021538305
<i>TYMS</i>	1.146977658	1.026076189	1.282124819	0.015825538

NSCLC, non-small cell lung cancer.

Table 5 The independent prognostic factors of NSCLC patients

Gene	coef	HR	HR.95L	HR.95H	P
<i>HIST1H2BH</i>	-0.220239949	0.802326258	0.729774307	0.882091103	5.25E-06
<i>PLK1</i>	0.303398284	1.354453815	1.199191723	1.529818043	1.04E-06

NSCLC, non-small cell lung cancer.


**Figure 6** The overexpression levels of HIST1H2BH and PLK1 were correlated with the gender and smoking history of LUAD patients in the UALCAN database. (A) HIST1H2BH; (B) PLK1. *, $P < 0.05$; **, $P < 0.01$; ***, $P < 0.001$. LUAD, lung adenocarcinoma.

Table 6 The value of risk model factors in LUAD

Clinical features	Type	HIST1H2BH	PLK1
Gender	Male vs. female	4.84E-03	1.47E-02
Age	Age (21–40 years) vs. age (81–100 years)	1.59E-02	NS
	Age (41–60 years) vs. age (61–80 years)	NS	1.91E-02
	Age (41–60 years) vs. age (81–100 years)	1.04E-04	NS
	Age (61–80 years) vs. age (81–100 years)	4.98E-03	NS
Smoking	Non smoker vs. smoker	4.77E-04	1.89E-05
	Non smoker vs. reformed smoker2	4.60E-03	1.86E-03
	Smoker vs. reformed smoker1	3.08E-03	2.84E-08
	Reformed smoker1 vs. reformed smoker2	2.58E-02	1.50E-06
Clinical stage	Stage1 vs. Stage3	NS	4.15E-02
	Stage1 vs. Stage4	NS	9.74E-03
Tissue type	NOS vs. mixed	7.16E-03	2.22E-03
	NOS vs. LBC-NonMucinous	3.45E-07	1.10E-07
	NOS vs. LBC-Mucinous	NS	5.93E-10
	NOS vs. mucinous carcinoma	2.41E-04	2.18E-02
	NOS vs. papillary	9.73E-07	NS
	NOS vs. micropapillary	7.48E-12	NS
	Mixed vs. LBC-NonMucinous	4.96E-03	2.56E-03
	Mixed vs. SolidPatternPredominant	1.84E-03	4.67E-02
	Mixed vs. LBC-Mucinous	NS	2.09E-07
	Mixed vs. papillary	1.25E-02	NS
	Mixed vs. micropapillary	4.82E-07	NS
	LBC-NonMucinous vs. SolidPatternPredominant	NS	9.71E-05
	LBC-NonMucinous vs. acinar	NS	2.24E-02
	LBC-NonMucinous vs. LBC-Mucinous	NS	4.67E-02
	SolidPatternPredominant vs. LBC-Mucinous	NS	1.51E-02
	SolidPatternPredominant vs. mucinous carcinoma	NS	1.48E-02
	SolidPatternPredominant vs. micropapillary	4.57E-02	NS
	Acinar vs. LBC-Mucinous	NS	8.57E-04
	LBC-Mucinous vs. mucinous carcinoma	NS	2.79E-02
	LBC-Mucinous vs. papillary	NS	1.09E-02
TP53 Mutant	TP53-Mutant vs. TP53-NonMutant	NS	9.91E-12

LUAD, lung adenocarcinoma.

Table 7 The value of risk model factors in LUSC

Clinical features	Type	HIST1H2BH	PLK1
LUSC	Normal vs. primary	<1E-12	<1E-12
Gender	Male vs. female	NS	4.31E-02
Age	Age (41–60 years) vs. age (61–80 years)	NS	1.41E-02
	Age (41–60 years) vs. age (81–100 years)	NS	2.34E-03
Smoking	Non smoker vs. smoker	1.84E-05	NS
	Non smoker vs. reformed smoker1	1.04E-03	NS
	Non smoker vs. reformed smoker2	3.15E-06	NS
	Smoker vs. reformed smoker1	NS	1.02E-02
	Reformed smoker1 vs. reformed smoker2	NS	3.68E-02
Clinical stage	Stage1 vs. Stage3	NS	2.90E-02
	Stage3 vs. Stage4	NS	2.70E-02
	NOS vs. papillary	1.68E-03	NS
Lymph node metastasis	N0 vs. N1	NS	6.36E-03
	N0 vs. N2	NS	1.32E-02
TP53 Mutant	TP53-Mutant vs. TP53-NonMutant	NS	2.68E-03

LUSC, lung squamous cell carcinoma.

suggested a poor prognosis for LUAD and LUSC patients, and the HR was 1.21 and 1.05, respectively. However, there was no statistical significance between the overexpression of PLK1 and the prognosis of LUSC patients (Figure S4B). In addition, we also found that HIST1H2BH and PLK1 had significant mutations in NSCLC (Figure S5).

Evaluation of the risk model in the prognosis of NSCLC patients

K-M survival analysis showed that high-risk NSCLC patients in the risk model had poor prognoses (Figure 9A). Univariate Cox regression analysis showed that the clinical stage, T stage, lymph node metastasis, and risk score were the influencing factors for the poor prognosis of NSCLC patients (Figure 9B). Multivariate Cox regression analysis showed that the clinical stage, T stage, and risk score were independent factors influencing the poor prognosis of NSCLC patients (Figure 9C). The nomogram showed that the clinical stage was the biggest influence on the survival of NSCLC patients, and the prognosis of NSCLC patients became worse as the stage increased. The second biggest influence was the risk score. As the risk score increased, the prognosis of NSCLC patients became worse (Figure 10).

The signaling mechanisms involved in the risk model

The high-risk group had significant roles in the cell cycle, mismatch repair, progesterone mediated oocyte maturation, oocyte meiosis, DNA replication, homologous recombination, ubiquitin-mediated proteolysis, proteasome, base excision repair, basal transcription factors, RNA degradation, p53 signaling pathway, glyoxylate and dicarboxylate metabolism, citrate cycle TCA cycle, lysine degradation, purine metabolism, NSCLC, and other cell types and functions (Figure 11 and Table 8).

The expression levels of HIST1H2BH and PLK1 were correlated with the expression level of PD-1

In lung tissue, the expression level of PLK1 was significantly correlated with the expression level of PD-1 (Figure S6A). Subgroup analysis showed that the expression level of PLK1 in normal tissues derived from LUAD and LUSC was significantly correlated with the expression level of PD-1 (Figure S6B,C). In LUAD and LUSC tissues, the expression level of PLK1 was significantly correlated with the expression level of PD-1 (Figure S6D,E). In lung tissue, the expression level of HIST1H2BH was significantly




Figure 7 Elevated expression levels of HIST1H2BH and PLK1 were related to the prognosis of NSCLC patients. (A,B,C) The OS, FP, and PPS of HIST1H2BH overexpression level in NSCLC patients; (D,E,F) the OS, FP, and PPS of PLK1 overexpression level in NSCLC patients. NSCLC, non-small cell lung cancer; OS, overall survival; PE, progression-free survival; PPS, post-progression survival.

correlated with the expression level of PD-1 (Figure S6F). Subgroup analysis showed that the expression level of HIST1H2BH in normal tissues derived from LUSC was significantly correlated with the expression level of PD-1 (Figure S6G). In LUSC tissue, the expression level of HIST1H2BH was significantly correlated with the expression level of PD-1 (Figure S6H).

Discussion

Recent years, NSCLC was one of the top tumors in the world in terms of morbidity and mortality (14). There were a variety of treatments for NSCLC patients that could improve the prognosis of NSCLC patients. However, the treatment effect was still not satisfactory. In recent years, immunotherapy was considered to be one of the effective

ways to improve advanced cancer. PD-1 inhibitor nivolumab could block the binding of PD-1 and PD-L1, promote T cells to participate in the recognition and killing of tumor cells, and prolong the survival of NSCLC patients (15). Therefore, this study explored the signaling mechanisms of PD-1 in the treatment of NSCLC and screen potential new target molecules for NSCLC progression to improve the long-term prognosis of NSCLC patients. In our current research, 48 PD-1 target molecules were involved in the cell division, cell proliferation, DNA replication, cell cycle, Lung Cancer, etc. studies indicated that cell proliferation, DNA replication, cell cycle played an important role in the development of NSCLC (16-18). For example, compared with paired normal lung tissues and bronchial epithelial cell, the expression level of TRIM13 in NSCLC tissues and cells was decreased. TRIM13 overexpression




Figure 8 HIST1H2B and PLK1 overexpression levels were related to the prognosis of lung cancer via the meta-analysis. (A,B) HIST1H2B and PLK1 expression levels; (C,D) the prognosis values of HIST1H2B and PLK1.




Figure 9 The risk model was related to the prognosis of NSCLC patients. (A) K-M survival analysis; (B) univariate Cox regression analysis; (C) multivariate Cox regression analysis. NSCLC, non-small cell lung cancer; K-M, Kaplan-Meier.




Figure 10 Construction of the risk model nomogram.




Figure 11 The signaling mechanisms involved in the risk model. (A) Cell cycle; (B) oocyte meiosis; (C) DNA replication; (D) homologous recombination; (E) ubiquitin mediated proteolysis; (F) basal transcription factors; (G) RNA degradation; (H) p53 signaling pathway; (I) pathway in cancer.

Table 8 The signaling mechanisms involved in the high-risk group

Name	Size	NES	NOM P
CELL_CYCLE	124	2.3562934	0
MISMATCH_REPAIR	23	2.2620559	0
PROGESTERONE_MEDIATED_OOCYTE_MATURATION	85	2.2228281	0
OOCYTE_MEIOSIS	112	2.2174144	0
DNA_REPLICATION	36	2.209795	0
ONE_CARBON_POOL_BY_FOLATE	17	2.2025287	0
NUCLEOTIDE_EXCISION_REPAIR	44	2.1026423	0
HOMOLOGOUS_RECOMBINATION	28	2.09409	0
PATHOGENIC_ESCHERICHIA_COLI_INFECTION	55	2.0320096	0
PYRIMIDINE_METABOLISM	97	1.9931018	0
UBIQUITIN_MEDIATED_PROTEOLYSIS	133	1.9535059	0
CHRONIC_MYELOID_LEUKEMIA	73	1.909263	0.001972387
AMINOACYL_TRNA_BIOSYNTHESIS	22	1.9242635	0.001992032
PANCREATIC_CANCER	70	1.8818412	0.002008032
BASAL_TRANSCRIPTION_FACTORS	35	1.8217084	0.004008016
PROTEASOME	44	1.9331573	0.004040404
BASE_EXCISION_REPAIR	33	1.8571749	0.004166667
PENTOSE_PHOSPHATE_PATHWAY	27	1.8049825	0.007722008
P53_SIGNALING_PATHWAY	68	1.7388005	0.007843138
RNA_DEGRADATION	57	1.7536411	0.008
SPLICEOSOME	126	1.7790519	0.008064516
GLYOXYLATE_AND_DICARBOXYLATE_METABOLISM	16	1.7177893	0.011904762
BLADDER_CANCER	42	1.5846695	0.016194332
PATHWAYS_IN_CANCER	325	1.5147824	0.016666668
GLIOMA	65	1.6071352	0.018108651
CITRATE_CYCLE_TCA_CYCLE	30	1.7157203	0.025590552
DEGRADATION	44	1.6497922	0.026859503
PURINE_METABOLISM	156	1.5934625	0.027139874
GAP_JUNCTION	89	1.5358167	0.027600849
PROSTATE_CANCER	89	1.5185701	0.030120483
DRUG_METABOLISM_OTHER_ENZYMES	51	1.5393608	0.031067962
THYROID_CANCER	29	1.5549417	0.032467533
SMALL_CELL_LUNG_CANCER	84	1.5911384	0.034836065
FRUCTOSE_AND_MANNOSE_METABOLISM	32	1.5649511	0.035363458
NON_SMALL_CELL_LUNG_CANCER	54	1.5211327	0.036885247
CYSTEINE_AND_METHIONINE_METABOLISM	34	1.4945567	0.03807615
AMYOTROPHIC_LATERAL_SCLEROSIS_ALS	52	1.4684905	0.038306452
N_GLYCAN_BIOSYNTHESIS	46	1.5310353	0.04761905
FC_GAMMA_R_MEDIATED_PHAGOCYTOSIS	95	1.5192246	0.04897959

NES, Normalized Enrichment Score; NOM P, nominal P value.

level in NCI-H1975 and SPC-A-1 cells could hinder cell proliferation, induce cell apoptosis, and inhibit tumor growth in nude mice (16). MSI1 could promote the proliferation and glucose metabolism of NSCLC cells and mediate the sensitivity to chemotherapy drugs in NSCLC cells. MSI1 could regulate protein kinase B (Akt) signaling activity (17).

Studies have shown that CCNA2 was the downstream target of miR-219-5p, and those were negatively correlated in esophageal squamous cell carcinoma (ESCC) tissues. Interference with CCNA2 expression could enhance the inhibitory effect of miR-219-5p on cell proliferation and the cell cycle (19). PLK1 inhibitor BI-6727 could induce mitotic mutations and DSB repair after radiation, thereby increasing the radiosensitivity of NSCLC cells. PLK1 silenced NSCLC cells showed that the gene expression related to DNA damage, replication, and repair were changed, including DNA-dependent protein kinase (DNAPK) and topoisomerase II α (TOPO2A) (18). The expression level of miR-223-5p in NSCLC tissues and cells was significantly down-regulated. The expression level of miR-223-5p was negatively correlated with the malignant degree of NSCLC. Overexpression of miR-223-5p could inhibit the proliferation, migration, and invasion of NSCLC cells *in vitro* and *in vivo*. This was related to the targeted regulation of E2F8 expression by miR-223-5p (19). Our study used Univariate Cox regression analysis to show that CCNA2, CHEK1, DLGAP5, E2F8, FOXM1, HIST1H2BH, HJURP, MKI67, PLK1, TPX2, and TYMS were the prognostic factors for NSCLC patients. Multivariate Cox regression analysis and AIC screening showed that HIST1H2BH and PLK1 were independent factors influencing the prognosis of NSCLC patients. It has been confirmed that some influencing factors played an important role in suppressing or promoting cancer in the progression of NSCLC (19-21), but there was still a further step to subdivide the influencing factors that need our follow-up confirmation. In addition, the risk model factors HIST1H2BH and PLK1 were overexpressed in LUAD and LUSC tissues and were related to the poor prognosis of patients. Survival analysis showed that high-risk NSCLC patients had a poor prognosis. Cox regression analysis showed that the clinical stage, T stage, and risk score were independent factors affecting the poor prognosis of NSCLC patients. The nomogram showed that as the risk score increases, the prognosis of NSCLC patients becomes worse. These results suggest that our risk model and nomogram have some value in predicting the prognosis

of NSCLC patients.

PD-1 has been used to treat advanced cancer patients in recent years and had certain therapeutic effects (1-3). In normal lung and cancer tissues, we found that there was a correlation between the factors of the constructed risk model and the expression level of PD-1. In lung tissue, the expression level of PLK1 was significantly correlated with the expression level of PD-1. The expression level of PLK1 in normal tissues derived from LUAD and LUSC was significantly correlated with the expression level of PD-1. In LUAD and LUSC tissues, the expression level of PLK1 was significantly correlated with the expression level of PD-1. In lung tissue, the expression level of HIST1H2BH was significantly correlated with the expression level of PD-1. The expression level of HIST1H2BH in normal tissues derived from LUSC was significantly correlated with the expression level of PD-1. In LUSC tissues, the expression level of HIST1H2BH was significantly correlated with the expression level of PD-1. It further showed that HIST1H2BH and PLK1 played an important role in the progression of NSCLC.

This study analyzed the gene expression levels in the blood of PD-1 inhibitor in the treatment of NSCLC, and verified it in the TCGA database to construct the risk model and nomogram for assessing the prognosis of NSCLC. PD-1 inhibitor expression could become a potential biomarker for cancer treatment, employing the risk model and nomogram as a guide to the patient's prognosis. The risk model factors we constructed have been verified by the multiple databases used in this study to have important clinical value, and multiple literature reported that PLK1 played an important role in the progression of NSCLC (22-24). However, this still needs further verification and will be part of our ongoing work. In general, PD-1 target molecules HIST1H2BH and PLK1 were overexpressed in LUAD and LUSC tissues. The increased expression levels of HIST1H2BH and PLK1 were related to the OS and PFS of NSCLC patients and were independent factors influencing the prognosis of NSCLC patients. In the risk model, high-risk NSCLC patients had a poor prognosis and were an independent factor influencing the poor prognosis of NSCLC patients. The high-risk group was enriched in signaling mechanisms such as cell cycle, DNA replication, homologous recombination, etc. The risk model based on PD-1 target molecules was helpful to assess the prognosis of NSCLC patients, and independent factors such as HIST1H2BH and PLK1 might become important prognostic biomarkers of NSCLC patients.

Acknowledgments

Funding: Health Commission of Hubei Province Scientific Research Project (WJ2019Q014) and Science and Technology Development Foundation of Xiaoping Chen (CXPJJH12000002-20200).

Footnote

Reporting Checklist: The authors have completed the REMARK reporting checklist. Available at <https://dx.doi.org/10.21037/atm-21-2963>

Conflicts of Interest: All authors have completed the ICMJE uniform disclosure form (available at <https://dx.doi.org/10.21037/atm-21-2963>). The authors have no conflicts of interest to declare.

Ethical Statement: The authors are accountable for all aspects of the work in ensuring that questions related to the accuracy or integrity of any part of the work are appropriately investigated and resolved. The study was conducted in accordance with the Declaration of Helsinki (as revised in 2013).

Open Access Statement: This is an Open Access article distributed in accordance with the Creative Commons Attribution-NonCommercial-NoDerivs 4.0 International License (CC BY-NC-ND 4.0), which permits the non-commercial replication and distribution of the article with the strict proviso that no changes or edits are made and the original work is properly cited (including links to both the formal publication through the relevant DOI and the license). See: <https://creativecommons.org/licenses/by-nc-nd/4.0/>.

References

1. Pu N, Gao S, Yin H, et al. Cell-intrinsic PD-1 promotes proliferation in pancreatic cancer by targeting CYR61/CTGF via the hippo pathway. *Cancer Lett* 2019;460:42-53.
2. Wang B, Qin L, Ren M, et al. Effects of Combination of Anti-CTLA-4 and Anti-PD-1 on Gastric Cancer Cells Proliferation, Apoptosis and Metastasis. *Cell Physiol Biochem* 2018;49:260-70.
3. Andrieu GP, Shafran JS, Smith CL, et al. BET protein targeting suppresses the PD-1/PD-L1 pathway in triple-negative breast cancer and elicits anti-tumor immune response. *Cancer Lett* 2019;465:45-58.
4. Xia L, Huang H, Xiao H, et al. Utilization of combined PD-L1 expression and neutrophil-to-lymphocyte ratio prior to surgery as a prognostic factor in non-small cell lung cancer with brain metastasis. *Transl Cancer Res* 2019;8:2864-77.
5. Wang J, Zhang R, Lin Z, et al. CDK7 inhibitor THZ1 enhances antiPD-1 therapy efficacy via the p38 α /MYC/PD-L1 signaling in non-small cell lung cancer. *J Hematol Oncol* 2020;13:99.
6. Ascierto PA, Del Vecchio M, Mandalá M, et al. Adjuvant nivolumab versus ipilimumab in resected stage IIIB-C and stage IV melanoma (CheckMate 238): 4-year results from a multicentre, double-blind, randomised, controlled, phase 3 trial. *Lancet Oncol* 2020;21:1465-77.
7. Czarnecka AM, Rutkowski P. An update on the safety of nivolumab for the treatment of advanced melanoma. *Expert Opin Drug Saf* 2020;19:409-21.
8. Fujimoto D, Yomota M, Sekine A, et al. Nivolumab for advanced non-small cell lung cancer patients with mild idiopathic interstitial pneumonia: A multicenter, open-label single-arm phase II trial. *Lung Cancer* 2019;134:274-8.
9. Morita R, Okishio K, Shimizu J, et al. Real-world effectiveness and safety of nivolumab in patients with non-small cell lung cancer: A multicenter retrospective observational study in Japan. *Lung Cancer* 2020;140:8-18.
10. Sugawara S, Lee JS, Kang JH, et al. Nivolumab with carboplatin, paclitaxel, and bevacizumab for first-line treatment of advanced non-squamous non-small cell lung cancer. *Ann Oncol* 2021; undefined; undefined.
11. Hoffner B, Leighl NB, Davies M. Toxicity management with combination chemotherapy and programmed death 1/programmed death ligand 1 inhibitor therapy in advanced lung cancer. *Cancer Treat Rev* 2020; undefined; 101979.
12. Hatae R, Chamoto K, Kim YH, et al. Combination of host immune metabolic biomarkers for the PD-1 blockade cancer immunotherapy. *JCI Insight* 2020;5:e133501.
13. Guo Q, Ke XX, Fang SX, et al. PAQR3 Inhibits Non-small Cell Lung Cancer Growth by Regulating the NF- κ B/p53/Bax Axis. *Front Cell Dev Biol* 2020; 8: 581919.
14. Siegel RL, Miller KD, Jemal A. Cancer statistics, 2020. *CA Cancer J Clin* 2020;70:7-30.
15. Schouten RD, Egberink L, Muller M, et al. Nivolumab in pre-treated advanced non-small cell lung cancer: long term follow up data from the Dutch expanded access program and routine clinical care. *Transl Lung Cancer Res* 2020;9:1736-48.
16. Xu L, Wu Q, Zhou X, et al. TRIM13 inhibited cell

- proliferation and induced cell apoptosis by regulating NF- κ B pathway in non-small-cell lung carcinoma cells. *Gene* 2019;715:144015.
17. Lang Y, Kong X, He C, et al. Musashi1 Promotes Non-Small Cell Lung Carcinoma Malignancy and Chemoresistance via Activating the Akt Signaling Pathway. *Cell Physiol Biochem* 2017;44:455-66.
 18. Ni Z, Yao C, Zhu X, et al. Ailanthone inhibits non-small cell lung cancer cell growth through repressing DNA replication via downregulating RPA1. *Br J Cancer* 2017;117:1621-30.
 19. Ma Q. MiR-219-5p suppresses cell proliferation and cell cycle progression in esophageal squamous cell carcinoma by targeting CCNA2. *Cell Mol Biol Lett* 2019;24:4.
 20. Yao D, Gu P, Wang Y, et al. Inhibiting polo-like kinase 1 enhances radiosensitization via modulating DNA repair proteins in non-small-cell lung cancer. *Biochem Cell Biol* 2018;96:317-25.
 21. Dou L, Han K, Xiao M, et al. miR-223-5p Suppresses Tumor Growth and Metastasis in Non-Small Cell Lung Cancer by Targeting E2F8. *Oncol Res* 2019;27:261-8.
 22. Yan L, Zhang Y, Li K, et al. miR-593-5p inhibit cell proliferation by targeting PLK1 in non small cell lung cancer cells. *Pathol Res Pract* 2020;216:152786.
 23. Shin SB, Jang HR, Xu R, et al. Active PLK1-driven metastasis is amplified by TGF- β signaling that forms a positive feedback loop in non-small cell lung cancer. *Oncogene* 2020;39:767-85.
 24. Reda M, Ngamcherdtrakul W, Gu S, et al. PLK1 and EGFR targeted nanoparticle as a radiation sensitizer for non-small cell lung cancer. *Cancer Lett* 2019;467:9-18.
- (English Language Editors: M. Bucci and J. Chapnick)

Cite this article as: Zhang YQ, Yuan Y, Zhang J, Lin CY, Guo JL, Liu HS, Guo Q. Evaluation of the roles and regulatory mechanisms of PD-1 target molecules in NSCLC progression. *Ann Transl Med* 2021;9(14):1168. doi: 10.21037/atm-21-2963




Figure S1 Disease-related to PD-1 target molecules. PD-1, programmed cell death protein 1.




Figure S2 The relationship between transcription factors and PD-1 target molecules. PD-1, programmed cell death protein 1.




Figure S3 HIST1H2B and PLK1 were unusual expression in LUAD and LUSC via the meta-analysis. (A) HIST1H2B; (B) PLK1. LUAD, lung adenocarcinoma; LUSC, lung squamous cell carcinoma.




Figure S4 HIST1H2B and PLK1 expression level were related to the OS in LUAD and LUSC tissues via the metaanalysis. (A) HIST1H2B; (B) PLK1. OS, overall survival; LUAD, lung adenocarcinoma; LUSC, lung squamous cell carcinoma.




Figure S5 HIST1H2BH and PLK1 had significant mutations in NSCLC. NSCLC, non-small cell lung cancer.




Figure S6 The expression levels of HIST1H2BH and PLK1 were correlated with PD-1 expression level. PD-1, programmed cell death protein 1.

Table S1 The biological functions of PD-1 target molecules via the GO

Category	Term	Count	P
BP	GO:0006334~nucleosome assembly	11	7.15E-13
BP	GO:0007067~mitotic nuclear division	13	2.06E-12
BP	GO:0007059~chromosome segregation	8	6.14E-10
BP	GO:0051301~cell division	11	2.85E-08
BP	GO:0008283~cell proliferation	11	4.34E-08
BP	GO:0034080~CENP-A containing nucleosome assembly	5	5.62E-06
BP	GO:0007062~sister chromatid cohesion	6	8.84E-06
BP	GO:0000086~G2/M transition of mitotic cell cycle	6	3.52E-05
BP	GO:0002227~innate immune response in mucosa	4	4.24E-05
BP	GO:0006260~DNA replication	6	6.35E-05
BP	GO:0000281~mitotic cytokinesis	4	6.69E-05
BP	GO:0019731~antibacterial humoral response	4	2.36E-04
BP	GO:0000910~cytokinesis	4	3.05E-04
BP	GO:0051726~regulation of cell cycle	5	3.64E-04
BP	GO:0051310~metaphase plate congression	3	4.76E-04
BP	GO:0031572~G2 DNA damage checkpoint	3	0.001351749
BP	GO:0050830~defense response to Gram-positive bacterium	4	0.001623959
BP	GO:0000070~mitotic sister chromatid segregation	3	0.002115826
BP	GO:0001578~microtubule bundle formation	3	0.002115826
BP	GO:0000732~strand displacement	3	0.002288156
BP	GO:0000082~G1/S transition of mitotic cell cycle	4	0.002732956
BP	GO:0007049~cell cycle	5	0.002894716
BP	GO:0007052~mitotic spindle organization	3	0.003041358
BP	GO:0000731~DNA synthesis involved in DNA repair	3	0.004123996
BP	GO:1901796~regulation of signal transduction by p53 class mediator	4	0.004733673
BP	GO:0000079~regulation of cyclin-dependent protein serine/threonine kinase activity	3	0.005100354
BP	GO:0046602~regulation of mitotic centrosome separation	2	0.010913626
BP	GO:0007018~microtubule-based movement	3	0.020737582
BP	GO:0030071~regulation of mitotic metaphase/anaphase transition	2	0.024391909
BP	GO:0006281~DNA repair	4	0.026418072
BP	GO:0006268~DNA unwinding involved in DNA replication	2	0.027065923
BP	GO:0001833~inner cell mass cell proliferation	2	0.032392459
BP	GO:0051382~kinetochore assembly	2	0.032392459
BP	GO:0010569~regulation of double-strand break repair via homologous recombination	2	0.042960087

Table S1 (continued)

Table S1 (continued)

Category	Term	Count	P
BP	GO:0006978~DNA damage response, signal transduction by p53 class mediator resulting in transcription of p21 class mediator	2	0.042960087
CC	GO:0005654~nucleoplasm	34	1.21E-17
CC	GO:0000788~nuclear nucleosome	9	1.44E-13
CC	GO:0000786~nucleosome	9	8.22E-11
CC	GO:0005634~nucleus	35	7.53E-10
CC	GO:0005819~spindle	8	2.06E-08
CC	GO:0000775~chromosome, centromeric region	6	3.11E-07
CC	GO:0005737~cytoplasm	30	1.27E-06
CC	GO:0000777~condensed chromosome kinetochore	6	2.59E-06
CC	GO:0000922~spindle pole	6	7.87E-06
CC	GO:0005876~spindle microtubule	4	1.85E-04
CC	GO:0015630~microtubule cytoskeleton	5	3.90E-04
CC	GO:0005813~centrosome	7	6.69E-04
CC	GO:0005874~microtubule	6	0.00108309
CC	GO:0000776~kinetochore	4	0.001118872
CC	GO:0000785~chromatin	4	0.001468433
CC	GO:0030496~midbody	4	0.004212549
CC	GO:0000784~nuclear chromosome, telomeric region	4	0.004304501
CC	GO:0005730~nucleolus	8	0.005348436
CC	GO:0000942~condensed nuclear chromosome outer kinetochore	2	0.010059237
CC	GO:0005829~cytosol	16	0.013417391
CC	GO:0005694~chromosome	3	0.02886921
CC	GO:0051233~spindle midzone	2	0.046907317
MF	GO:0046982~protein heterodimerization activity	11	2.45E-07
MF	GO:0003677~DNA binding	18	4.03E-07
MF	GO:0019901~protein kinase binding	9	4.94E-06
MF	GO:0008017~microtubule binding	7	1.56E-05
MF	GO:0005515~protein binding	36	1.49E-04
MF	GO:0005524~ATP binding	12	0.001252552
MF	GO:0042393~histone binding	4	0.003930198
MF	GO:0008022~protein C-terminus binding	4	0.011807477
MF	GO:0003777~microtubule motor activity	3	0.018448317
MF	GO:0042826~histone deacetylase binding	3	0.029009124

PD-1, programmed cell death protein 1; GO, gene ontology; BP, biological processes; CC, cellular component; MF, molecular function.