Study on the interaction mechanism between *C-reactive protein* and platelets in the development of acute myocardial infarction

Yubao Liu¹, Shuhui Lai², Lijie Liang³, Donghai Zhang¹

¹Department of Intensive Care Medicine, The Second Affiliated Hospital of Qiqihar Medical College, Qiqihar, China; ²The First Clinical Medical College of Nanchang University, Nanchang, China; ³Ultrasound Department, The Second Affiliated Hospital of Qiqihar Medical College, Qiqihar, China

Contributions: (I) Conception and design: Y Liu, D Zhang; (II) Administrative support: L Liang; (III) Provision of study materials or patients: Y Liu, D Zhang, L Liang; (IV) Collection and assembly of data: Y Liu, L Liang; (V) Data analysis and interpretation: D Zhang, Y Liu; (VI) Manuscript writing: All authors; (VII) Final approval of manuscript: All authors.

Correspondence to: Yubao Liu. Department of Intensive Care Medicine, The Second Affiliated Hospital of Qiqihar Medical College, No. 37, West Zhonghua Road, Jianhua District, Qiqihar 161006, China. Email: liuyubao0452@163.com.

Background: Myocardial infarction (MI) is the single most critical event in coronary disease. Platelets are involved in the processes of acute MI (AMI). They lack nuclear DNA but retain megakaryocyte mRNAs, hence, their transcriptome could provide information preceding coronary events. However, their mechanisms are not clear. In this study, we obtained a gene expression atlas of platelets from patients after their very first AMI, and our purpose was to clarify the mechanisms of platelet involvement in the occurrence of AMI through bioinformatics analyses and animal models of AMI *in vivo*.

Methods: We obtained a gene expression atlas of platelets from patients after their very first AMI from the Gene Expression Omnibus (GEO). Differentially expressed genes (DEGs) were retrieved using R language. Weighted gene co-expression network analysis (WGCNA) was implemented in order to construct a gene co-expression correlation network among DEGs. Animal models of AMI *in vivo* were constructed to confirm the results of the bioinformatics analysis.

Results: Gene integration analysis yielded 2,852 DEGs ($P < 0.05$, $|\log_2FC| > 1$). Bioinformatics analysis demonstrated a significant association between *C-reactive protein* (CRP) and *Staphylococcus aureus* infection (SAI) ($P = 0.015$). Data from *in vivo* experiments showed that CRP increased significantly in AMI rats ($P < 0.001$), and the expression of *FCGR2B* mRNA and *HLA-DRB4* mRNA was elevated in response to the increase of CRP ($P < 0.001$).

Conclusions: From the results of this study, we speculate that in the development of AMI, the increase in CRP activates platelets and induces platelets to play an anti-inflammatory role.

Keywords: Acute myocardial infarction (AMI); *C-reactive protein* (CRP); platelets

Submitted Apr 30, 2021. Accepted for publication Jun 16, 2021.

doi: 10.21037/atm-21-2733

View this article at: <https://dx.doi.org/10.21037/atm-21-2733>

Introduction

Acute myocardial infarction (AMI) is a disease in which the heart experiences a sudden deprivation of circulating blood, inducing acute and constant ischemia and hypoxia of the heart and increasing the risk of death (1). Proximately every 40 seconds, someone in the United States experiences

a myocardial infarction (MI). AMI accounts for roughly 80% of patients in cardiogenic shock (CS) (2). AMI has two identified subtypes: ST-segment elevation MI (STEMI) and non-STEMI (3). STEMI generally presents as constant ischemic chest pain and elevated serum myocardial necrosis markers, alongside the typical ST-segment elevations on

electrocardiogram (ECG) (4). It accounts for approximately 25–40% of AMI (5). The majority of patients with STEMI can receive percutaneous coronary intervention and drugs to relieve symptoms, but it remains an impossible task to rescue the necrotic and apoptotic myocardial cells (6). Unfortunately, no significant decrease has been observed in the risk of death of AMI patients in hospital over the past decades, yet the number of affected patients in China is expected to increase to 23 million by the year 2030 (7). Even if timely treatment exists, acute myocardial ischemic injury and ensuing ischemia-reperfusion injury continue to be challenging issues. Furthermore, radiological and other auxiliary examination approaches, as well as elucidating applicable biomarkers, are regarded as highly necessary for the early diagnosis of this event (8).

Platelets play a key role in thrombotic processes that limit the patency of the recanalized, infarct-related coronary artery and contribute to reperfusion injury (9). The vascular importance of platelets has been attributed to their essential role in mediating MI (10). Studies have shown that platelets become activated during MI (11), but the mechanism of their participation in AMI is not clear. Thus, in our study, we aim to clarify the mechanisms of platelet involvement in the occurrence of AMI through bioinformatics analyses and animal models of AMI *in vivo*. We present the following article in accordance with the ARRIVE reporting checklist (available at <https://dx.doi.org/10.21037/atm-21-2733>).

Methods

Microarray-based analysis concerning up-regulated differentially expressed genes (DEGs)

The study was conducted in accordance with the Declaration of Helsinki (as revised in 2013).

First, the GSE24519 dataset was retrieved from the Gene Expression Omnibus (GEO) database (available at <http://www.ncbi.nlm.nih.gov/geo/>), which comprised of platform files and original files (CEL files). Next, background correction was implemented, whereupon quantile normalization, prosummarization, and log₂ conversion along with a missing values supplement related to the matrix data of GEO were followed with the use of the Robust Multi-array Average algorithm with the R “affy” and “impute” packages. Up-regulated DEGs of GSE24519 were identified and matrices were constructed using the R “limma” package with $|\log_2FC| > 1$, alongside P value < 0.05 working as the threshold.

Weighted gene co-expression network analysis (WGCNA) construction

The GSE24519 dataset included 17 AMI patients with overall survival details, and thus qualified for WGCNA construction. Next, the module eigengenes were linked with the clinical data of AMI patients. A data matrix concerning gene expression in GSE24519 was then constructed, with the first 25% of genes in disease samples exhibiting the largest variance chosen to serve as the input dataset applied for the follow-up WGCNA with the use of the R “WGCNA” package. The outlier sample was then detected and excluded by employing the sample hierarchical clustering method, which was prior to the selection of a qualified soft-threshold power to obtain a standard scale-free network. Thereafter, by means of adjacency, topological overlap matrix (TOM) construction, and corresponding dissimilarity calculation, a gene dendrogram was generated and module identification was carried out with the use of dynamic tree cut, with a minimum module size of 50. Subsequently, module eigengenes cluster analysis was carried out to merge extremely similar modules (with dissimilarity less than 0.25), followed by calculating the correlation of the module eigengene with the AMI clinical phenotype. The module was then selected and used for further analyses, then gene significance (GS) and module membership (MM) were acquired and used for identifying the modules linked to the clinical traits of AMI.

Enrichment analysis

In the gene network in accordance with the scale-free distribution, genes exhibiting identical expression would be modulated together, linked functionally or by shared pathways. Upon topological overlap calculation, genes in the aforementioned chosen module showing a positive association with *C-reactive protein (CRP)* were investigated. Subsequently, Gene Ontology (GO) and Kyoto Encyclopedia of Genes and Genomes (KEGG) analyses were performed via WebGestalt (available at <http://www.webgestalt.org/>). Functions presenting with $P < 0.05$ were considered statistically significant.

Animals and animal models

Forty purchased specific-pathogen free (SPF) Sprague-Dawley rats (180–220 g, 6–8 W, Hunan, China) were randomized into four groups, namely the AMI group,

control group, *CRP* group, and sham operation group. Prior to the experiment, 2% pentobarbital sodium, sterile equipment packages, and a 5-F bipolar temporary pacing electrode were prepared. The procured rats were food-fasted for a duration of 12 hours, water-fasted for a duration of 6 hours, partially depilated on the back, and weighed. The rats in the AMI group were anesthetized with 2% pentobarbital sodium (0.3 mL/100 g), fixed in the supine position, and intubated with air intake maintained with a ventilator. The leads of the animal ECG monitoring machine were inserted into the subcutaneous muscle layer of limbs according to the operation requirements. After obtaining stable readings on the ECG, second left intercostal thoracotomy was performed to expose the pericardium, and the coronary artery was completely ligated with surgical suture 1–2 mm from the left anterior descending branch. After a few seconds, the ECG showed STEMI, the heart was thereby reset, and the chest was closed layer by layer. The rats in the *CRP* group were fed a conventional diet and injected with *CRP* (recombinant human *CRP*) through the tail vein (3 mg/kg). Meanwhile, the rats randomized into the sham operation group underwent puncture in the left anterior descending branch of the left coronary artery without ligation, and the rest of the process was the same as the model establishment. The rats in the control group were fed a conventional diet without any treatment. Experiments were approved by ethics board of The Second Affiliated Hospital of Qiqihar Medical University, in compliance with Qiqihar Medical University guidelines for the care and use of animals.

Preparation of plasma

Platelet-rich plasma (PRP) is defined as autologous serum with a high concentration of platelets and growth factors, which is obtained by centrifugation of whole blood (11). All rats were anesthetized with 2% pentobarbital sodium at a dose of 0.3 mL/100 g, then blood samples were obtained from the abdominal aorta and collected in a test tube containing 2 mL heparin (5,000 IU/mL) and 800 μ L 10% sodium citrate, with an average of 6 mL blood collected per animal. PRP can be prepared via one-step or two-step centrifugation procedures (12). The obtained PRP and platelet-poor plasma (PPP) were prepared and stored at 20 °C. According to the protocol, 10% CaCl_2 (10 μ L/200 μ L PRP) was added to activate PRP before use (13).

Enzyme-linked immunosorbent assay (ELISA)

ELISA is amongst the most significant biochemical analysis techniques for diagnosing and monitoring a plethora of diseases (14). The content of *CRP* in PPP was detected as per the manufacturer's instructions of the rat high sensitivity *CRP* ELISA kit (Sigma, USA).

Real-time quantitative polymerase chain reaction (RT-qPCR)

The total RNA content of PRP, which was isolated using the TRIzol reagent, underwent cDNA reverse transcription with the use of Reverse Transcription kits (PN4366596, acquired from Applied Biosystems). Expression patterns of long non-coding RNAs (lncRNAs) in tissues were quantified using KAPA SYBR FAST qPCR kits (KK4601, acquired from Kapa Biosystems). *GAPDH* was employed as the internal control, and levels of expressed genes were assessed with the $2^{-\Delta\Delta C_t}$ method combined with MxPro 4.00 (Stratagene), with primer sequences shown in Table 1.

Statistical analysis

All data involved in this investigation were expressed as mean $\pm$ standard deviation (mean $\pm$ SD), with a minimum of three repeated independent experiments. Data were analyzed using Graphpad prism 5.0 software. Two-group data comparisons were implemented with Student's *t*-test. $P < 0.05$ indicated statistical significance.

Results

Identification of DEGs in AMI

The GSE24519 dataset included 17 AMI patients with detailed gene expression profile information. GSM604591, GSM604585, and GSM604589 were excluded from subsequent analysis in GSE24519 as outlier samples. Therefore, a total of 14 samples with gene expression profile information were included in this study, and 2,852 DEGs ($P < 0.05$, $|\log_2 \text{FC}| > 1$) were identified.

Construction of WGCNA and identification of the key module

WGCNA is a well-recognized systematic biological technique often applied for analyzing the expression of

Table 1 The used primer sequences

Gene name	Primer sequence	
	Forward sequence (5'-3')	Reverse sequence (5'-3')
<i>FCGR2B</i>	GTTGGGGCTGAGAACAACAAAT	ACAGGGAGCTTCAGGACTCA
<i>HLA-DRB4</i>	AGACAAGTTCACCCACACAG	AGCATCAAACCTCCAGTGCT
<i>GAPDH</i>	TGTGTCCGTCGTGGATCTGA	CCTGCTTCACCACCTTCTTGA

genes in varied samples. It is capable of forming either a cluster or module that contains genes presenting with identical expression (15). Once genes are present in a module, they exhibit a tendency to possess similar biological functions, and the associations of modules with sample features can be investigated (16).

The GSM604591, GSM604585, and GSM604589 datasets were removed from follow-up analyses in GSE24519 as outlier samples, leaving 14 samples with survival data that were included in WGCNA (Figure 1A,B). The power of $\beta=6$ (scale free $R^2=0.9$) was chosen as the soft-threshold in order to construct a scale-free network (depicted in Figure 1C,D). Thirteen modules related to gene co-expression were finally identified following removal of the grey module with the use of merged dynamic tree cutting (Figure 1E). A heat map depicting the TOM of the 1000 selected genes showed that each module worked as an independent verification to one another (depicted in Figure 1F). Subsequently, we identified the green-yellow module, representing a key module showing the most significant association with CRP ($R^2=0.86$, $P=7e-05$; Figure 1G). Next, we attempted to draw scatter plots concerning GS relative to MM in the green-yellow module with GS for CRP (correlation =0.84, $P=3.5e-19$; Figure 1H). Gene expression profiles from platelet samples in The Cancer Genome Atlas (TCGA) identified that these 68 genes were over-expressed in AMI samples (Figure 1I).

Genes showing a positive association with CRP are uncovered to be enriched in Staphylococcus aureus infection (SAI)

We then investigated the possible biological functions of CRP-related genes in AMI. The results of GO analysis demonstrated that these genes were primarily enriched in biological regulation, metabolic process, cell communication, membrane, nucleus, protein binding, and ion binding, among others (Figure 2A). Further KEGG analysis yielded the correlation between CRP and SAI (Figure 2B).

Model diagnosis

The successful establishment of AMI model can be judged by ECG, serum biomarkers or echocardiography, and each method has its important diagnostic significance. In this study, the method of observing the animal signs and ECG in the process of model replication was used as the standard to detect the successful establishment of AMI model. The results of this experiment showed that after ligation of the left anterior descending coronary artery, the myocardial tissue of the left ventricular anterior wall became pale and contraction weakened. Physiological signal acquisition and processing system software system shows that the ECG is converted from normal QRS waveform to typical STEMI. These results indicate that the AMI model was successfully established.

CRP increased in AMI rats

After 6 hours in the animal model, the average levels of CRP in the normal group, sham operation group, AMI group, and CRP group were $1,049\pm162$, $1,674\pm190$, $6,577\pm836$, and $7,949\pm910$ ng/mL, respectively. These findings showed that the content of CRP was increased in the AMI group in contrast to the sham group (**, $P<0.01$), indicating that AMI can precipitate an increase in CRP. In contrast to the controls, the level of CRP was even higher in the CRP group (**, $P<0.01$) (Figure 3A,B). Meanwhile, when compared with the normal group, the CRP level in the sham operation group was much higher, which was considered to be related to surgical trauma. However, the increased amplitude and trend were even lower than those in the AMI and CRP groups.

High expression of FCGR2B and HLA-DRB4 links to CRP

In order to verify whether the increase of CRP can

Figure 1 Development of the weighted co-expression network. (A,B) Hierarchical cluster dendrogram of samples acquired from GSE24519. (C,D) Scale-free fit indicator along with mean connectivity related to varied soft-threshold powers. $\beta=6$ for scale-free topology test. (E) Hierarchical cluster dendrogram of differential genes based on topological overlap. In this figure, modules are indicative of the branches in the cluster tree. (F) A heat map depicting the TOM of the selected 1,000 genes. In this figure, darker color reflects higher overlap while lighter color reflects lower overlap. Gene dendrogram along with module assignment is exhibited along the left and top. (G) Correlation assessment of module eigengenes with clinical traits. Columns reflect clinical traits and each row corresponds to a module eigengene. Every cell contains correlation and P values, with the green-yellow module containing *CRP*. (H) Scatter plots depicting GS relative to MM in the green-yellow module with GS for *CRP* (correlation = 0.84, $P=3.5e-19$). (I) The expression matrix of 68 DEGs from platelet samples in TCGA database. In this figure, blue reflects lower expression, red reflects higher expression, and white reflects no expression difference in the genes. TOM, topological overlap matrix; GS, gene significance; MM, module membership; *CRP*, *C-reactive protein*; DEG, differentially expressed gene; TCGA, The Cancer Genome Atlas.

Figure 2 Genes displaying a positive association with *CRP* were found to be enriched in SAI. (A) GO analysis of genes displaying a positive association with *CRP*. (B) KEGG analysis of genes displaying positive relation with *CRP*. *CRP*, *C-reactive protein*; SAI, *Staphylococcus aureus* infection; GO, Gene Ontology; KEGG, Kyoto Encyclopedia of Genes and Genomes.

influence the expression levels of infection-related genes, the expression patterns of *FCGR2B* and *HLA-DRB4* in the four groups were detected using RT-qPCR. The results illustrated increased mRNA expression levels of *FCGR2B* and *HLA-DRB4* in the AMI group in contrast to the sham operation group (***, $P < 0.001$) (Figure 4). Meanwhile, the mRNA expression levels of *FCGR2B* and *HLA-DRB4* in the CRP group were increased in relation to the control group (***, $P < 0.001$). The increase of *CRP* may affect the infection-related pathway and aggravate the disease process.

Discussion

AMI, along with heart failure that often ensues, are major causes of morbidity and mortality worldwide (17). Interestingly, the work of our peers has shown that inflammation is key to the pathogenesis of atherosclerosis and acute coronary events (18). Inflammatory responses are a common occurrence after AMI, and can worsen ischemia-reperfusion injuries, triggering augmented infarct size and worse prognoses (19). Moreover, studies have

Figure 3 The average levels of *CRP* in the normal group, sham operation group, AMI group, and *CRP* group after 6 hours in the animal model. (A) Standard curve of *CRP*. (B) Statistical chart of *CRP*. ***, $P < 0.001$. *CRP*, C-reactive protein; AMI, acute myocardial infarction; OD, optical density.

also documented persistently elevated levels of *CRP*, an inflammation biomarker, in patients following AMI (20).

Firstly, findings obtained in our study from the differential analysis of the GSE24519 dataset using WGCNA revealed a total of 2,852 DEGs. Subsequent findings indicated that the green-yellow module was highly positively correlated with *CRP* traits. Meanwhile, *in vivo* experimentation findings demonstrated that the AMI group exhibited significantly higher levels of *CRP* in contrast to the sham operation group in the STEMI model rats ($P < 0.001$). Elevated levels of *CRP* have been previously documented in numerous conditions such as renal failure, thyroid disease, and congestive heart failure (21). In our study, GO and KEGG analyses of the 68 genes in the *CRP*-related green-yellow module in AMI demonstrated that these genes were primarily involved in biological regulation, metabolic process, stimulation response, cell communication, and other biological processes. KEGG pathway analysis illustrated that there were significant differences in three pathways, including SAI, hematopoietic

cell lineage, and glycosphingolipid biosynthesis ($P < 0.05$), wherein SAI was found to be the most significant pathway ($P = 0.0148$). The correlation between AMI and infection was first acknowledged during the early 20th century in the context of influenza epidemics. More recently, the association between AMI and SAI has been identified by a case-control and self-control design investigation, which is in accordance with our findings (22).

Inflammatory processes after the advent of MI attract predominant attention in the field of cardiovascular research (23). AMI patients who present with constantly elevated biomarkers related to inflammation (20), in particular *CRP*, have been reported to be at a greatly increased risk of developing further cardiovascular events (24). *CRP* is an evolutionarily conserved protein, with homologs present in vertebrates and numerous invertebrates (25). Elevated *CRP* levels are a common occurrence in inflammatory conditions, cardiovascular diseases, and infections (26). The plasma concentration of *CRP* was previously shown to deviate by at least 25% in the context

Figure 4 The mRNA expression levels of *FCGR2B* and *HLA-DRB4* in the normal group, sham operation group, AMI group, and CRP group after 6 hours in the animal model. ***, $P < 0.001$. AMI, acute myocardial infarction; CRP, C-reactive protein.

of inflammatory disorders (27). Traditionally, CRP works as a non-specific marker related to inflammation. However, research in the last decade has further highlighted the direct involvement of CRP in inflammation and atherosclerosis, as well as other cardiovascular diseases. More importantly, one particular study highlighted CRP as the most powerful predictor and risk factor for cardiovascular disease (28). Furthermore, CRP levels in humans within 6 hours after coronary artery occlusion are known to reflect the inflammatory state of myocardial ischemia (29). The inflammatory reaction is activated within 4–8 hours, and presents with gradual increases in CRP levels (30). In this study, after 6 hours of modeling, CRP in mice increased significantly.

Another notable finding in our study was the increased expression levels of *FCGR2B* and *HLA-DRB4* in addition to CRP in AMI. Inherently, *FCGR2B* and *HLA-DRB4* serve as susceptible genes in autoimmune diseases (31) and play critical roles in maintaining the homeostasis of the immune response (32). *FCγRs* are well-recognized cell surface receptors that are known to participate in a variety of autoimmune diseases by recognizing the anti-IgG antibody and mediating the immune response, and interacting with the *Fc* domain of IgG. Currently, there are three different classes of *FcγR* recognized in humans: *FcγRII* (CD32), *FcγRI* (CD64), and *FcγRIII* (CD16) (33). Platelets adhere and aggregate around bacteria, typically utilizing their *GP2b/3a*, *FcγRIIIa*, and IgG receptors in conjunction with fibrinogen or fibronectin (34). Various researchers have further documented significant correlations between *HLA-DRB* and

CRP in patients with rheumatoid arthritis, and these genes are even involved in the immune process of myocarditis (35,36). In this study, we established a STEMI rat model *in vivo*, and assessed the expression patterns of the AMI-related genes *FCGR2B* and *HLA-DRB4* in the SAI pathway using RT-PCR. Our findings showed that the expression levels of *FCGR2B mRNA* and *HLA-DRB4 mRNA* in PRP in the AMI group were significantly elevated in contrast to the sham operation group. Compared with the normal group, the expression levels of *FCGR2B mRNA* and *HLA-DRB4 mRNA* in PRP in the CRP group were also significantly elevated.

Studies have reported that platelets are involved in the inflammatory response (10). Combined with the results of this study, we speculate that in the development of AMI, the increase in CRP activates platelets and induces platelets to play an anti-inflammatory role.

Acknowledgments

Funding: This study was funded by Qiqihar Science and Technology Program (SFGG-201731).

Footnote

Reporting Checklist: The authors have completed the ARRIVE reporting checklist. Available at <https://dx.doi.org/10.21037/atm-21-2733>

Data Sharing Statement: Available at <https://dx.doi.org/10.21037/atm-21-2733>

[org/10.21037/atm-21-2733](https://doi.org/10.21037/atm-21-2733)

Conflicts of Interest: All authors have completed the ICMJE uniform disclosure form (available at <https://dx.doi.org/10.21037/atm-21-2733>). The authors have no conflicts of interest to declare.

Ethical Statement: The authors are accountable for all aspects of the work in ensuring that questions related to the accuracy or integrity of any part of the work are appropriately investigated and resolved. Experiments were approved by ethics board of The Second Affiliated Hospital of Qiqihar Medical University, in compliance with Qiqihar Medical University guidelines for the care and use of animals. The study was conducted in accordance with the Declaration of Helsinki (as revised in 2013).

Open Access Statement: This is an Open Access article distributed in accordance with the Creative Commons Attribution-NonCommercial-NoDerivs 4.0 International License (CC BY-NC-ND 4.0), which permits the non-commercial replication and distribution of the article with the strict proviso that no changes or edits are made and the original work is properly cited (including links to both the formal publication through the relevant DOI and the license). See: <https://creativecommons.org/licenses/by-nc-nd/4.0/>.

References

- Lewis EF, Li Y, Pfeffer MA, et al. Impact of cardiovascular events on change in quality of life and utilities in patients after myocardial infarction: a VALIANT study (valsartan in acute myocardial infarction). *JACC Heart Fail* 2014;2:159-65.
- Lavall MC, Bagatini MD, Thomé GR, et al. Extracellular hydrolysis of adenine nucleotides and nucleoside adenosine is higher in patients with ST elevation than non-ST elevation in acute myocardial infarction. *Clin Lab* 2015;61:761-7.
- Kapur NK, Thayer KL, Zweck E. Cardiogenic shock in the setting of acute myocardial infarction. *Methodist Debakey Cardiovasc J* 2020;16:16-21.
- Gnavi R, Ruscioni R, Dalmaso M, et al. Gender, socioeconomic position, revascularization procedures and mortality in patients presenting with STEMI and NSTEMI in the era of primary PCI. Differences or inequities? *Int J Cardiol* 2014;176:724-30.
- Kook HY, Jeong MH, Oh S, et al. Current trend of acute myocardial infarction in Korea (from the Korea Acute Myocardial Infarction Registry from 2006 to 2013). *Am J Cardiol* 2014;114:1817-22.
- Kloner RA, Dai W, Hale SL, et al. Approaches to improving cardiac structure and function during and after an acute myocardial infarction: acute and chronic phases. *J Cardiovasc Pharmacol Ther* 2016;21:363-7.
- Li J, Li X, Wang Q, et al. ST-segment elevation myocardial infarction in China from 2001 to 2011 (the China PEACE-Retrospective Acute Myocardial Infarction Study): a retrospective analysis of hospital data. *Lancet* 2015;385:441-51. Retracted and republished in: *Lancet* 2015;385:402.
- Ye XD, He Y, Wang S, et al. Heart-type fatty acid binding protein (H-FABP) as a biomarker for acute myocardial injury and long-term post-ischemic prognosis. *Acta Pharmacol Sin* 2018;39:1155-63.
- Gawaz M, Neumann FJ, Ott I, et al. Platelet function in acute myocardial infarction treated with direct angioplasty. *Circulation* 1996;93:229-37.
- Koupenova M, Clancy L, Corkrey HA, et al. Circulating platelets as mediators of immunity, inflammation, and thrombosis. *Circ Res* 2018;122:337-51.
- Xiao SJ, Zhou YF, Wu Q, et al. Uncovering the differentially expressed genes and pathways involved in the progression of stable coronary artery disease to acute myocardial infarction using bioinformatics analysis. *Eur Rev Med Pharmacol Sci* 2021;25:301-2.
- Yazigi Junior JA, Dos Santos JB, Xavier BR, et al. Quantification of platelets obtained by different centrifugation protocols in SHR rats. *Rev Bras Ortop* 2015;50:729-38.
- Xu J, Gou L, Zhang P, et al. Platelet-rich plasma and regenerative dentistry. *Aust Dent J* 2020;65:131-42.
- Zhdanov A, Keefe J, Franco-Waite L, et al. Mobile phone based ELISA (MELISA). *Biosens Bioelectron* 2018;103:138-42.
- Beckerman P, Qiu C, Park J, et al. Human kidney tubule-specific gene expression based dissection of chronic kidney disease traits. *EBioMedicine* 2017;24:267-76.
- Tang J, Kong D, Cui Q, et al. Prognostic genes of breast cancer identified by gene co-expression network analysis. *Front Oncol* 2018;8:374.
- Ong SB, Hernández-Reséndiz S, Crespo-Avilan GE, et al. Inflammation following acute myocardial infarction: multiple players, dynamic roles, and novel therapeutic opportunities. *Pharmacol Ther* 2018;186:73-87.
- Newby LK. Inflammation as a treatment target after acute

- myocardial infarction. *N Engl J Med* 2019;381:2562-3.
19. Akodad M, Sicard P, Fauconnier J, et al. Colchicine and myocardial infarction: a review. *Arch Cardiovasc Dis* 2020;113:652-9.
 20. Thackeray JT, Hupe HC, Wang Y, et al. Myocardial inflammation predicts remodeling and neuroinflammation after myocardial infarction. *J Am Coll Cardiol* 2018;71:263-75.
 21. Liu Z, Ma C, Gu J, et al. Potential biomarkers of acute myocardial infarction based on weighted gene co-expression network analysis. *Biomed Eng Online* 2019;18:9.
 22. McNamara JF, Harris PNA, Chatfield MD, et al. Acute Myocardial Infarction and Community-Acquired Staphylococcus aureus Bloodstream Infection: an observational cohort study. *Clin Infect Dis* 2020. [Epub ahead of print]. doi: 10.1093/cid/ciaa1197.
 23. Rischpler C. Acute myocardial infarction. *Q J Nucl Med Mol Imaging* 2016;60:236-51.
 24. Hennessy T, Soh L, Bowman M, et al. The Low Dose Colchicine after Myocardial Infarction (LoDoCo-MI) study: a pilot randomized placebo controlled trial of colchicine following acute myocardial infarction. *Am Heart J* 2019;215:62-9.
 25. Black S, Kushner I, Samols D. C-reactive protein. *J Biol Chem* 2004;279:48487-90.
 26. Du Clos TW, Mold C. C-reactive protein: an activator of innate immunity and a modulator of adaptive immunity. *Immunol Res* 2004;30:261-77.
 27. Sproston NR, Ashworth JJ. Role of C-reactive protein at sites of inflammation and infection. *Front Immunol* 2018;9:754.
 28. Osimo EF, Baxter LJ, Lewis G, et al. Prevalence of low-grade inflammation in depression: a systematic review and meta-analysis of CRP levels. *Psychol Med* 2019;49:1958-70.
 29. Geluk CA, Post WJ, Hillege HL, et al. C-reactive protein and angiographic characteristics of stable and unstable coronary artery disease: data from the prospective PREVENT cohort. *Atherosclerosis* 2008;196:372-82.
 30. Anzai T. Inflammatory mechanisms of cardiovascular remodeling. *Circ J* 2018;82:629-35.
 31. Yuksel M, Xiao X, Tai N, et al. The induction of autoimmune hepatitis in the human leucocyte antigen-DR4 non-obese diabetic mice autoimmune hepatitis mouse model. *Clin Exp Immunol* 2016;186:164-76.
 32. Su K, Wu J, Edberg JC, et al. A promoter haplotype of the immunoreceptor tyrosine-based inhibitory motif-bearing FcγRIIb alters receptor expression and associates with autoimmunity. I. Regulatory FCGR2B polymorphisms and their association with systemic lupus erythematosus. *J Immunol* 2004;172:7186-91.
 33. Vogelpoel LT, Baeten DL, de Jong EC, et al. Control of cytokine production by human fc gamma receptors: implications for pathogen defense and autoimmunity. *Front Immunol* 2015;6:79.
 34. Loughman A, Fitzgerald JR, Brennan MP, et al. Roles for fibrinogen, immunoglobulin and complement in platelet activation promoted by Staphylococcus aureus clumping factor A. *Mol Microbiol* 2005;57:804-18.
 35. Kliment B, Nefic H, Prodanovic N, et al. Association of biomarkers of inflammation and HLA-DRB1 gene locus with risk of developing rheumatoid arthritis in females. *Rheumatol Int* 2019;39:2147-57.
 36. Şelli ME, Thomas AC, Wraith DC, et al. A humanized HLA-DR4 mouse model for autoimmune myocarditis. *J Mol Cell Cardiol* 2017;107:22-6.
- (English Language Editor: C. Betlazar-Maseh)

Cite this article as: Liu Y, Lai S, Liang L, Zhang D. Study on the interaction mechanism between *C-reactive protein* and platelets in the development of acute myocardial infarction. *Ann Transl Med* 2021;9(12):1012. doi: 10.21037/atm-21-2733